

Target-specific requirements for RNA interference can arise through restricted RNA amplification despite the lack of specialized pathways

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

Reviewed preprint version 1

May 28, 2024 (this version)

Sent for peer review

March 18, 2024

Posted to preprint server

March 1, 2024

Daphne R. Knudsen-Palmer, Pravrutha Raman, Farida Etefa, Laura De Ravin, Antony M. Jose 

Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, USA. Biological Sciences Graduate Program, University of Maryland, College Park, USA

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Since double-stranded RNA (dsRNA) is effective for silencing a wide variety of genes, all genes are typically considered equivalent targets for such RNA interference (RNAi). Yet, loss of some regulators of RNAi in the nematode *C. elegans* can selectively impair the silencing of some genes. Here we show that such selective requirements can be explained by an intersecting network of regulators acting on genes with differences in their RNA metabolism. In this network, the Maelstrom domain-containing protein RDE-10, the intrinsically disordered protein MUT-16, and the Argonaute protein NRDE-3 work together so that any two are required for silencing one somatic gene, but each is singly required for silencing another somatic gene, where only the requirement for NRDE-3 can be overcome by enhanced dsRNA processing. Quantitative models and their exploratory simulations led us to find that (1) changing *cis*-regulatory elements of the target gene can reduce the dependence on NRDE-3, (2) animals can recover from silencing in non-dividing cells and (3) cleavage and tailing of mRNAs with UG dinucleotides, which makes them templates for amplifying small RNAs, is enriched within 'pUG zones' matching the dsRNA. Similar crosstalk between pathways and restricted amplification could result in apparently selective silencing by endogenous RNAs.

eLife assessment

This **valuable** study shows how an intersecting network of regulators acting on genes with differences in their RNA metabolism explains why the loss of some regulators of RNAi in *C. elegans* can selectively impair the silencing of some target genes. The evidence presented is **convincing**, as the authors use a combination of computational modeling and RNAi assays to support their conclusions.

<https://doi.org/10.7554/eLife.97487.1.sa2>

Introduction

Double-stranded RNA (dsRNA) can trigger the conserved mechanism of RNA interference (RNAi) to degrade mRNA of matching sequence (Fire et al., 1998), and thus silence gene expression, in many organisms. This conservation has made dsRNA-based drugs useful in crops (Das and Sherif, 2020), insects (Vogel et al., 2019), and humans (Zhu et al., 2022). While a dsRNA-based drug can be designed using just the mRNA sequence of any target gene, the intracellular effectiveness of the drug and the ease with which an organism could escape the drug by developing resistance are difficult to predict. Predicting both efficacy and susceptibility to resistance for each target could inform the selection of a suitable target from two or more equivalent candidates. Extensive characterization of RNAi in the nematode *C. elegans* (reviewed in (Seroussi et al., 2022)) makes it a suitable system to examine how differences between target genes and reliance on specific regulators contribute to efficacy and resistance.

A skeletal pathway that is required for gene silencing in response to the addition of dsRNA has been worked out in *C. elegans* (Fig. 1A). Long dsRNA is imported through the transmembrane protein SID-1 (Feinberg and Hunter, 2003; Winston et al., 2002), after which it is bound by the dsRNA-binding protein RDE-4 (Tabara et al., 2002), which recruits the endonuclease DCR-1 (Knight and Bass, 2001) to cleave the long dsRNA into smaller dsRNAs (Parker et al., 2006). The primary Argonaute protein RDE-1 (Parrish and Fire, 2001; Tabara et al., 1999) cleaves one strand of the smaller dsRNA (Steiner et al., 2009) and associates with the other, making it a 1° short interfering RNA (siRNA) that can guide the recognition of target mRNAs of matching sequence (siRNAs; processing, pink). After recognition by RDE-1-bound siRNAs, the target mRNAs are cleaved and the 5' fragments are stabilized through the addition of 3' UG-dinucleotide repeats (Preston et al., 2019) by the nucleotidyltransferase RDE-3 (Chen et al., 2005) to form pUG RNAs (Shukla et al., 2020), which act as templates for the amplification of 2° siRNAs (Pak and Fire, 2007) by RNA-dependent RNA polymerases. This amplification of silencing signals through the production of 2° siRNAs is facilitated by the intrinsically disordered protein MUT-16 (Phillips et al., 2012; Zhang et al., 2011), the Maelstrom domain-containing protein RDE-10 (Yang et al., 2012; Zhang et al., 2012), and their interactors (Phillips et al., 2012; Uebel et al., 2018; Yang et al., 2012; Zhang et al., 2012). These 2° siRNAs are bound by one of several Argonautes (Yigit et al., 2006), resulting in the eventual degradation of target mRNAs in the cytoplasm, which requires a cytoplasmic Argonaute, and/or co-transcriptional silencing of the target gene in the nucleus, which requires a nuclear Argonaute (e.g., NRDE-3 (Guang et al., 2008) in somatic cells). Although it is difficult to compare the silencing of two different genes by controlling all relevant variables, past studies have highlighted gene-specific differences in the efficacy of RNAi under different conditions (e.g., when RNAi is enhanced through the loss of the exonuclease ERI-1 (Kennedy et al., 2004), when nuclear silencing is blocked in somatic cells through loss of NRDE-3 (Raman et al., 2017), or when different concentrations of dsRNA are used (Zhuang and Hunter, 2011)). Understanding the sources of such differences and the underlying mechanisms will improve our ability to design efficacious dsRNA drugs that are difficult to evade through the development of resistance.

Here we analyze the requirements for silencing two exemplar genes and use quantitative modeling to advance a parsimonious view of RNAi in somatic cells. We show that MUT-16, RDE-10, and NRDE-3 are each required for the silencing of *bli-1*, but any two of these proteins are sufficient for *unc-22* silencing. These differences can be explained by differences in the thresholds for silencing the two genes using an intersecting network of regulators but not by parallel pathways of regulation after primary mRNA recognition. The requirement for NRDE-3 but not for MUT-16 or RDE-10 can be bypassed by enhancing the processing of dsRNA, suggesting that loss of NRDE-3 has the least impact on the efficiency of silencing. A dynamic model of RNA changes during silencing




Fig. 1.

RNA interference of two somatic targets show stark differences in their requirements for MUT-16 and RDE-10.

(A) Overview of RNA interference in somatic cells. Double-stranded RNA (dsRNA, blue) enters the cell through the importer SID-1 (import, teal), after which it is processed by the dsRNA-binding protein RDE-4 and the endonuclease Dicer into 1° short interfering RNAs (siRNAs) that are bound by the primary Argonaute RDE-1 (1° processing, pink). mRNA transcripts (green) recognized by these 1° siRNAs are modified after cleavage by the 3' addition of UG repeats (pUG RNA) and act as templates for the amplification of 2° siRNAs aided by the intrinsically disordered protein MUT-16, the Maelstrom-domain containing protein RDE-10, and RNA-dependent RNA polymerases (2° amplification, orange). These 2° siRNAs can bind secondary Argonaute(s) (e.g., NRDE-3), which can then recognize additional complementary targets (2° recognition) and cause gene silencing. See text for details. (B) Gene schematics depicting the mutant alleles found in a genetic screen (black) and/or created using genome editing (blue). Black boxes indicate exons and red dots indicate locations of mutations. Allele names (e.g., *jam139*) and expected amino acid change in the corresponding proteins (e.g., mutation of a tyrosine codon to a stop codon, Y294*) are indicated. See Fig. S1 for details of genetic screen. (C and D) Response to *bli-1* or *unc-22* RNAi in different mutants. For each mutant, the fraction of animals that showed *bli-1* silencing or *unc-22* silencing (fraction silenced) and the numbers of animals scored (n) are shown. Asterisks indicate $p < 0.05$ for each comparison (brackets) using Wilson's estimates with continuity correction and error bars represent 95% confidence interval. (C) Of five isolates with a mutation in *mut-16*, four (*jam138*, *jam140*, *jam141*, and *jam247*) failed to silence *bli-1* (blue) but retained *unc-22* silencing (orange). The other mutant failed to silence both genes and additionally had a mutation in *rde-10* (*mut-16(jam139) rde-10(jam248)*). (D) Mutants created using genome editing recapitulated the selective silencing of *unc-22* in *mut-16(-)* single mutants (*mut-16(jam148)*) and the failure to silence both genes in *mut-16(-) rde-10(-)* double mutants (*mut-16(jam148) rde-10(jam206)*). Using genome editing to recreate the *jam248* mutation, which is expected to make a mutant protein (RDE-10(S228F)) that disrupts the Maelstrom domain (see Fig. S1), resulted in animals (*rde-10(jam196)*) that showed *unc-22* silencing but not *bli-1* silencing. (E) Selective requirement for a regulator could reflect two underlying mechanisms of RNA silencing: (1) Two parallel pathways (left, A/B path vs C/D path) that are differentially used for different target genes; or (2) One intersecting network (right, A/B/C/D network) with quantitative contributions by all regulators along with different thresholds for each target gene.

by dsRNA reveals several criteria for efficient RNA silencing. Insights from modeling led us to discover the influence of *cis*-regulatory regions on the requirements for RNAi, the recovery of animals from RNAi within non-dividing cells, and a dearth of pUG RNA production by 2° siRNAs.

Results

Two genes with different thresholds for silencing reveal an intersecting network of regulators that mediate RNA interference

To identify regulators of RNA interference (RNAi), we performed a primary screen for mutants that disrupt the maintenance of mating-induced silencing of a transgene followed by a secondary screen for mutants that are also defective in the silencing of endogenous genes by ingested dsRNA (Fig. S1A). Mating males with a transgene that expresses fluorescent proteins to hermaphrodites that lack the transgene can initiate silencing in progeny that lasts in descendants for hundreds of generations (Devanapally et al., 2021 [↗](#)), providing a stable strain that can be mutagenized to look for mutations that result in the recovery of expression from the fluorescent transgene. Of the 15 fertile mutants that showed re-expression, whole-genome sequencing followed by *in silico* complementation (see Materials and Methods), revealed five mutants that had premature stop codons in *mut-16* (Fig. 1B [↗](#)), a known regulator of RNAi that is required for the production of secondary siRNAs (Phillips et al., 2012 [↗](#); Uebel et al., 2018 [↗](#); Zhang et al., 2011 [↗](#)). MUT-16 is detectable in the germline localized within perinuclear foci, but it is also found throughout the soma (Uebel et al., 2018 [↗](#)). MUT-16 is required for the silencing of all tested somatic targets except the muscle gene *unc-22*, which showed residual silencing ('+++ vs '+' but not '-' in (Zhang et al., 2011 [↗](#))) consistent with its early identification as a sensitive target for RNAi (Fire et al., 1998 [↗](#)). While all five putative *mut-16* mutants failed to silence the hypodermal gene *bli-1* (Fig. 1C [↗](#), left), only four of the five showed *unc-22* silencing (Fig. 1C [↗](#), right). Upon further analysis of the mutant that failed to silence *unc-22*, we found that this mutant also contained a missense mutation in RDE-10, another known regulator of RNAi that is required for the production of secondary siRNAs (Yang et al., 2012 [↗](#); Zhang et al., 2012 [↗](#)). This missense mutation (Ser228Phe) is expected to disrupt the Maelstrom domain of RDE-10 (Fig. S1C), and thus could result in a loss of RDE-10 function. To eliminate possible confounding effects of multiple mutations in strains isolated from a genetic screen, we used Cas9-mediated genome editing to introduce mutations in *mut-16* (null) and/or *rde-10* (null or a missense mutation that encodes Ser228Phe) in a wild-type background (Fig. 1B [↗](#)). While the newly created *mut-16*(null) mutants showed *unc-22* silencing as expected, *mut-16*(null) *rde-10*(null) (Fig. 1D [↗](#), right, Fig. S2B) double mutants failed to silence *unc-22*. Since *unc-22* is a particularly sensitive target for RNAi (Fire et al., 1998 [↗](#)), this lack of *unc-22* silencing in the absence of two regulators with roles in the amplification of 2° siRNAs suggests that 1° siRNA production and RDE-1-mediated recognition of the mRNA is likely not sufficient to cause silencing of most genes. These observations suggest that MUT-16 and RDE-10 are redundantly or additively required for silencing *unc-22* and that the Maelstrom domain of RDE-10 is required for this function. Since the primary Argonaute RDE-1 is required for the silencing of all somatic targets (Fig. 1A [↗](#); Parrish and Fire, 2001 [↗](#); Tabara et al., 1999 [↗](#)), including *unc-22*, we propose that MUT-16 and RDE-10 act in parallel downstream of RDE-1 to promote the amplification of 2° siRNA. This branching of the RNAi pathway downstream of RDE-1 could result in strictly parallel pathways where MUT-16 and RDE-10 are used to silence different sets of genes (Fig. 1E [↗](#), left) or in an intersecting network where both regulators contribute to the silencing of all genes (Fig. 1E [↗](#), right).

Additional observations suggest differences in the requirements for silencing *bli-1* and *unc-22*. Animals that lack MUT-16 (Fig. 1D [↗](#)), RDE-10 (Fig. 1D [↗](#)) or the somatic nuclear Argonaute NRDE-3 (Raman et al., 2017 [↗](#)) fail to silence *bli-1* but not *unc-22*. However, *rde-10*(-); *nrde-3*(-) double mutants fail to silence *unc-22* (Yang et al., 2012 [↗](#)). Therefore, if there were strictly parallel pathways downstream of MUT-16 and RDE-10, then NRDE-3 would be expected to function

downstream of MUT-16 but parallel to RDE-10. To test this possibility, we generated *nrde-3(-)* mutants using genome editing (Fig. 2A) and compared silencing in single mutants and double mutant combinations using the newly generated mutants lacking MUT-16, RDE-10, or NRDE-3. As expected, all single mutants failed to silence *bli-1* but silenced *unc-22*. Surprisingly, all double mutants failed to silence both *bli-1* and *unc-22* (Fig. 1D and 2B). This requirement for any two of MUT-16, RDE-10, or NRDE-3 suggests that the RNAi pathway cannot be strictly parallel downstream of RDE-1 (see Fig. S2A).

The stark differences in the extents of silencing *bli-1* (~0%) versus *unc-22* (~100%) (Fig. 1D and 2B) in animals lacking MUT-16, RDE-10, or NRDE-3 suggest that there could be target-specific pathways for silencing, tissue-specific differences in the expressions of RNA regulators, or more parsimoniously, that each regulator contributes to the silencing of both targets as part of an intersecting network, through transcriptional gene silencing (TGS) and/or post-transcriptional gene silencing (PTGS) (Fig. 2C), with *unc-22* being more sensitive to silencing than *bli-1*. For such an intersecting network with quantitative contributions by multiple regulators of RNAi to explain the silencing of somatic targets, including targets like *unc-22* and *bli-1* that show dramatic differences, it should be possible to identify values for the relative contributions of each regulatory path (N_m = from MUT-16 to NRDE-3, N_r = from RDE-10 to NRDE-3, O_m = from MUT-16 to other Argonautes, and O_r = from RDE-10 to other Argonautes in Fig. 2C, left) and for gene-specific thresholds (T_{bli-1} = level of BLI-1 function below which a defect is detectable, and T_{unc-22} = level of UNC-22 function below which a defect is detectable) that are consistent with all experimental data ('constraints' in Fig. 2C, right). Of the 100,000 sets of parameters simulated, 145 sets satisfied all experimental constraints (Fig. 2D). These allowed parameter sets were obtained despite the conservative assumption that the levels of mRNA knockdown for detecting observable defects for *bli-1* and *unc-22* are similar. Relaxing this assumption will lead to a larger number of allowed parameter sets. These valid parameter sets included cases with different relative contributions from RDE-10 and MUT-16 to NRDE-3-dependent silencing for a range of threshold differences for silencing *bli-1* versus *unc-22* (Fig. 2D, left). Furthermore, extreme contributions of MUT-16 versus RDE-10 (Fig. 2D, middle) or NRDE-3 versus other Argonautes (Fig. 2D, right) were absent. Finally, only thresholds for *bli-1* silencing that are less than ~5.5x the threshold for *unc-22* silencing were supported despite the allowed range of up to 100x (Fig. 2D). Consistent with different quantitative contributions to silencing by each regulator, reducing the availability of *unc-22* dsRNA revealed a graded silencing response: *mut-16(-) < rde-10(-) < nrde-3(-)* (Fig. 2E; ~5% silencing in *mut-16(jam148)* ~15% in *rde-10(jam206)* and ~70% in *nrde-3(jam205)*). Consistent with the possibility of differential contributions from each regulator for different targets (Fig. 2D), while partial silencing is observable in the absence of NRDE-3 but not MUT-16 or RDE-10 when the muscle gene *unc-54* is targeted (Fig. 2F, left), partial silencing is observed in the absence of MUT-16 but not NRDE-3 or RDE-10 when the hypodermal gene *dpy-7* is targeted (Fig. 2F, right).

Taken together, our results are consistent with a single network for RNAi targeting somatic genes where intersecting regulatory pathways downstream of mRNA recognition provide quantitative contributions to silencing.

The genetic requirement for NRDE-3, but not for MUT-16 and/or RDE-10, can be bypassed by enhancing dsRNA processing

The production of pUG RNAs and 2° siRNAs requires the participation of mRNA (Fig. 1A), making the contributions of some steps during RNAi gene-specific. Therefore, genes could differ in their dependence on proteins required for steps downstream of dsRNA processing and 1° siRNA production. Such differential dependencies could in principle be overcome by increasing the amount of available processed dsRNA and/or 1° siRNA when alternative parallel paths are available (e.g., loss of NRDE-3, MUT-16, or RDE-10 in Figs. 1 and 2) but not when no alternative paths are available (e.g., loss of both MUT-16 and RDE-10 in Fig. 2B) or when the




Fig. 2.

Gene-specific requirements and complex redundancy can arise from a single RNA regulatory network.

(A) Schematic (as in Fig. 1) depicting *nrde-3* alleles. (B) Feeding RNAi of *bli-1* and *unc-22*. Fractions silenced, numbers scored, comparisons, asterisks, and error bars are as in Fig. 1. Single mutants lacking NRDE-3 (*nrde-3(jam205)*) fail to silence *bli-1* but not *unc-22*. Double mutants fail to silence both targets. (C and D) Mutual constraints among parameters required for a single RNA regulatory network to support experimental results. (C, left) Model for a single network of interactors that regulate all RNAi targets in somatic cells. All targets require import (SID-1) and processing (RDE-4 and RDE-1) of dsRNA. Branching after 1st siRNA processing results in four distinct paths (*Nm*, *Nr*, *Om*, *Or*) that together contribute to gene silencing, which could occur through co-transcriptional gene silencing (TGS) and/or post-transcriptional gene silencing (PTGS) mechanisms. (C, right) Representation of simulation workflow. First, random values between 0 and 2 were drawn for each of the four variables (*Nm*, *Nr*, *Om*, *Or*). Second, constraints were added based on the experimental results in Fig. 2B and Fig. 1D. Third, allowed values that satisfied all experimental conditions were culled. Of 100,000 sets of random values simulated (0 to 2 for *Nm*, *Nr*, *Om*, *Or* and 0 to 100 for the ratio of thresholds T_{bli-1}/T_{unc-22}), 145 were consistent with all observed responses to RNAi. These allowed numbers reveal the domain of parameter values that support the observed range of gene silencing outcomes using feeding RNAi. (D, left) The contribution of NRDE-3 via MUT-16 (*Nm*) versus that via RDE-10 (*Nr*) for different ratios of thresholds for *bli-1* versus *unc-22* silencing (T_{bli-1}/T_{unc-22}) are shown. (D, center and right) The relative contributions to silencing that require MUT-16 (*Nm* + *Om*, D, center) or NRDE-3 (*Nm* + *Nr*, D, right) do not frequently take extreme values and both support a low value for the ratio of thresholds ($T_{bli-1}/T_{unc-22} < \sim 5.5$ despite allowed values of up to 100). (E) Feeding RNAi of *unc-22* assayed as in Fig. 1, but using aged plates, resulting in weaker RNAi. Animals that lack MUT-16 (*mut-16(jam148)*) have the most severe defect, followed by animals lacking RDE-10 (*rde-10(jam206)*), which is followed by animals lacking NRDE-3 (*nrde-3(jam205)*). (F) Feeding RNAi of *unc-54* or *dpy-7*. Fractions silenced, numbers scored, comparisons, asterisks, and error bars are as in Fig. 1. Silencing of *unc-54* showed a partial dependency on NRDE-3, while silencing of *dpy-7* showed a partial dependency on MUT-16, suggesting that the quantitative requirement for a regulator can differ depending on the target.

increase in dsRNA processing is insufficient. To test these predictions, we increased dsRNA processing and examined silencing in animals lacking different regulators required for the silencing of *bli-1* and/or *unc-22*.

One approach for increasing dsRNA processing is the release of factors such as the endonuclease DCR-1 from competing endogenous pathways by removing the exonuclease ERI-1 (Lee et al., 2006). In addition to the increased availability of DCR-1 when ERI-1 is removed, downstream factors involved in siRNA amplification (e.g. MUT-16, MUT-2/RDE-3, RDE-10/11, etc.) and 2° Argonautes (e.g. WAGOs) would be more available to contribute to silencing in response to ingested dsRNA. We used available *eri-1* mutants (Fig. 3A, *mg366*) and mutants generated using Cas9-mediated genome editing (Fig. 3A, *jam260* to *jam264*) to test if requirements for silencing *bli-1* and/or *unc-22* could be bypassed. Loss of ERI-1 enabled *bli-1* silencing in animals lacking NRDE-3, but not in animals lacking RDE-10 or MUT-16 (Fig. 3B). Furthermore, loss of *eri-1* was not sufficient for the complete rescue of *unc-22* silencing in animals lacking any two of these three regulators (Fig. 3C). An alternative approach for increasing dsRNA processing is the overexpression of the dsRNA-binding protein RDE-4, which recruits dsRNA for processing by DCR-1 (8, 10). Minimal amounts of RDE-4 can support RNAi as evidenced by silencing in *rde-4(-)* adult progeny of *rde-4(+/-)* hermaphrodites (Fig. S7E in (Marré et al., 2016)) and in *rde-4(-)* animals with trace levels of ectopic expression from multicopy *rde-4(+)* transgenes (Fig. 2 in (Raman et al., 2017)). We found that even hemizygous males expressing *rde-4(+)* from a single-copy transgene driving expression in the germline and the intestine under the control of the *mex-5* promoter (Marré et al., 2016) was sufficient for rescuing both *bli-1* and *unc-22* silencing (Fig. 3D). Similar expression of *rde-1(+)*, however, was not sufficient for rescuing silencing in *rde-1(-)* animals (Fig. 3D), suggesting that small amounts of RDE-4 but not RDE-1 are sufficient for RNAi. RDE-4 can be selectively overexpressed in the hypodermis using a single-copy transgene with a *nas-9* promoter (overexpression evident in Fig. S3; and selectivity demonstrated in Fig. 4C in (Raman et al., 2017)). This hypodermal expression of *rde-4(+)* was sufficient to enable *bli-1* silencing in an otherwise *rde-4(-); nrde-3(-)* background (Fig. 3E). Thus, either loss of ERI-1 or overexpression of RDE-4 can bypass the need for NRDE-3 for silencing *bli-1*, suggesting that the requirement for this regulator does not reflect a specific need for a particular regulator (NRDE-3) but reflects a larger amount of silencing signals required for reducing *bli-1* function sufficiently to cause a detectable defect. However, loss of ERI-1 and/or overexpression of RDE-4 could not compensate for the loss of RDE-10 or MUT-16 for *bli-1* silencing (Fig. 3F), suggesting that these regulators make a more substantial contribution to *bli-1* silencing than NRDE-3. These observations further support the idea that 1° siRNAs alone are not sufficient to cause silencing, consistent with the lack of *unc-22* silencing in *mut-16(-) rde-10(-)* double mutants (Fig. 1B, 1D). One explanation for these results is that in *eri-1(-); nrde-3(-)* double mutants (Fig. 3G), a different 2° Argonaute is able to compensate for the lack of NRDE-3, whereas in *mut-16(-); eri-1(-)* or *rde-10(-); eri-1(-)* double-mutants, the reduction of 2° siRNAs is too great to cause a detectable Bli-1 defect.

Taken together, these results suggest that gene-specific requirements for some proteins that function in RNAi do not reflect different pathways for silencing different genes, but rather a quantitative requirement for regulators acting as part of an intersecting RNA regulatory network.

Quantitative modeling of RNA interference and mRNA production provides rationales for a variety of target-specific outcomes

The many protein regulators of RNAi drive changes in RNA metabolism, including the production of new RNA species (1° siRNA, 2° siRNA, and pUG RNA), that are associated with the targeted gene. Although these changes can be indicators of RNA silencing, the quantitative relationship between such RNA intermediates and the extent of gene silencing measured as a reduction in function of the targeted gene or its mRNA levels is unclear. A priori, reduction in the mRNA levels of a gene could depend on universal processing of imported dsRNA, production of secondary small RNAs




Fig. 3.

Gene-specific requirements for NRDE-3 can be bypassed in two ways.

(A) Gene schematics (as in Fig. 1) of *rde-4* and *eri-1*. (B and C) Loss of ERI-1 can bypass the NRDE-3 requirements for silencing *bli-1* but not the other requirements for silencing *bli-1* or *unc-22*. Feeding RNAi targeting *bli-1* (B) or *unc-22* (C) with fractions silenced, numbers scored, comparisons, asterisks, and error bars as in Fig. 1. (B) Loss of ERI-1 (*mg366*, *jam263*, and *jam264* alleles) can compensate for the role of NRDE-3 (*tm1116* and *jam205* alleles) but not of RDE-10 (*jam206* allele) or MUT-16 (*jam148*, *jam265*, *jam266*, *jam267*, and *jam268* alleles) in *bli-1* silencing. See Table S3 for additional information. (C) Silencing of *unc-22* is not restored by loss of ERI-1 (*mg366*, *jam260*, *jam261*, and *jam262* alleles) in mutants that also lack any two of *mut-16* (*jam148* and *jam240* alleles), *rde-10* (*jam196* and *jam206* alleles), or *nrde-3* (*jam205* allele). See Table S3 for additional information. (D to E) Overexpression of RDE-4 in the hypodermis can bypass the requirement for NRDE-3 in *bli-1* silencing. (D) Minimal amounts of RDE-4 are sufficient for somatic silencing. (top) Schematic depicting generation of male progeny with paternal inheritance of a single-copy transgene (*Si[...]*) that expresses *rde-4(+)* or *rde-1(+)* under the control of the *mex-5* promoter (*mex-5p*) in the germline (green) of *rde-4(-)* or *rde-1(-)* animals, respectively (germline- and intestine-enriched RDE, based on rescue of RNAi in *rde-1(-)* animals (39)). (bottom) Male cross progeny with the transgene were scored after feeding only F1 animals, showing that unlike animals with germline- and intestine-enriched RDE-1, animals with similarly enriched RDE-4 can rescue both *unc-22* and *bli-1* silencing. Thus, small amounts of RDE-4 potentially mis-expressed in the hypodermis or a non-autonomous effect of RDE-4 from the germline or intestine is sufficient for silencing in the muscle and hypodermis. † indicates $p < 0.05$ when compared to either wild type or the *rde-4(-)* mutant and other symbols are as in (B). (E) Silencing of *bli-1* is restored in *nrde-3(tm1116); rde-4(ne301)* double mutants when *rde-4(+)* is overexpressed in the hypodermis (*Si[nas-9p::rde-4(+)]*). (F) Silencing of *bli-1* is not restored in animals lacking MUT-16 or RDE-10, despite the overexpression of RDE-4 in the hypodermis and/or the loss of ERI-1. (G) Summary depicting the bypass of NRDE-3 when ERI-1 is eliminated and/or RDE-4 is overexpressed. The increase in dsRNA processing increases the contributions of NRDE-3-independent paths to silencing.




Fig. 4.

A quantitative model allows exploration of parameters for RNAi in *C. elegans*.

(A) RNAi can be modeled to understand (from *left*) the responding populations of animals, the outline of mechanism, the process highlighting RNA species, the biochemical sub-steps (e.g., 2° siRNA production using pUG RNA, *right*). This study focuses on the overall RNA-mediated processes that accompany RNAi. (B) Schematic (*left*) and ordinary differential equations (*right*) describing the production and turnover of different RNA species. Rate constant for 1° siRNA processing from dsRNA (*k1*), binding constant for 1° siRNA binding target mRNA (*k2*), rate constant for pUG RNA production (*k3*), rate constant for 2° siRNA production (*k4*), binding constant for 2° siRNAs binding mRNA (*k5*) or pre-mRNA (*k6*), rate constant for export and splicing of transcripts from the nucleus (*k7*), rate constant for repression of transcription (*k8*), and rate constant for new transcript production (*k9*). Other terms are described in the figure. (C) Relative changes in the concentrations of each RNA ([RNA] vs. time in a.u.; dsRNA, mRNA, pre-mRNA, pUG RNA, 1° siRNA, and 2° siRNA) for an example set of parameters (all turnover rates = 0.05, *k1* = 1, *k2* = 0.01, *k3* = 1, *k4* = 0.05 * l_m = 0.5, *k5* = 0.01, *k6* = 0.01, *k7* = 0.1, *k8* = 0.05, *k9* = 7.5) are illustrated. A reduction to 10% of initial mRNA concentration is designated as the threshold for detecting a defect upon knockdown ([m]_{kd}), the time needed to reach the threshold (t_{kd}) and the time for which mRNA levels remain below the threshold (t_{kd}) are also indicated. (D) Relationship between the duration of knockdown and the time to knockdown (t_{kd} and t_{kd} are as in (C)). (E) Relationship between mRNA concentration and 2° siRNA accumulation. The minimum mRNA concentrations and maximum 2° siRNA concentrations reached for different transcripts with two different binding constants of 2° siRNAs binding to mRNA (*k5* = 0.1, red and *k5* = 0.01, blue) are plotted. Also see Fig. S5. (F) Impact of doubling transcription on transcripts with different knockdown parameters. Each transcript is colored based on its initial duration of knockdown (t_{kd}, blue to red gradient) before a 2-fold increase in the rate constant for transcription (*k9*) and the resultant fractional change in the duration of knockdown t_{kd} is plotted against that in the time to knockdown t_{kd}. (G) Genes with higher turnover are harder to knockdown. Response of mRNAs and their respective 1° siRNA with the same steady-state concentrations but with different rates of mRNA turnover (solid lines: T_m = 0, large dashes: T_m = 0.05, small dashes: T_m = 0.5) upon addition of 10 molecules of dsRNA are shown. (*inset*) Relationship of the minimum concentration of mRNA ([mRNA]_{min}) with its T_m in response to a fixed amount of dsRNA.

with the participation of gene-specific mRNAs, and downregulation of pre-mRNA and mRNA influenced by pre-existing gene-specific RNA metabolism. To understand how these gene-specific factors could influence RNA silencing, we began by analyzing the impact of a few characteristics of a gene on mRNA (m) and pre-mRNA levels (p) after RNAi. We first used a sequential equilibrium model, where we assume each step must be completed before beginning the next, e.g. dsRNAs are fully processed into 1° siRNAs, then 1° siRNAs can recognize transcripts to result in the production of pUG RNAs, and so on (Fig. S4A and Supplemental Methods). We tested parameters that would result in varying levels of target RNA knockdown (790 of 1 million simulated parameters resulted in $[m]_i < [m]$, $[m]_i > 0$, and $[p]_i > 0$; Fig. S4B). Under this simple model, we found that (1) RNAi can result in different residual concentrations of RNAs for different genes (Fig. S4C); (2) for a given gene, silencing can alter the ratio of pre-mRNA to mRNA (Fig. S4D and S4E); and (3) effective targeting of mRNA by primary or secondary small RNAs is required for strong silencing (Fig. S4F). These observations hint at the influence of gene-specific factors on the functional outcome of RNAi and impel the exploration of a more detailed dynamic model.

A qualitative outline of the molecular mechanism for RNAi in *C. elegans* has been deduced based on a variety of studies over the last two decades (mechanism outline; Fig. 4A [↗](#), left), but the cellular, subcellular, and kinetic details of every step remain obscure. Quantitative modeling of RNAi – or indeed any process of interest – could be done at many scales (Fig. 4A [↗](#), right) based on the level of understanding sought and experimental data available for testing predictions. For example, the responses of different animals in a population to dsRNA exposure (population model; Fig. 4A [↗](#), right) or the changes in key RNA species after entry of dsRNA into the cytosol (process model; Fig. 4A [↗](#), right) could be modeled. At yet greater detail, one step such as the amplification of small RNAs using pUG RNA templates could be modeled by incorporating sequence bias, processivity of RdRP, etc. (biochemical model; Fig. 4A [↗](#), right). Of these scales, we focused on the process model because early process models of RNAi (e.g., [Bergstrom et al., 2003 \[↗\]\(#\)](#)) were proposed before crucial discoveries on the biogenesis of 2° siRNAs without forming long dsRNA ([Pak and Fire, 2007 \[↗\]\(#\)](#)) and the stabilization of mRNA templates as pUG RNAs ([Preston et al., 2019 \[↗\]\(#\)](#); [Shukla et al., 2020 \[↗\]\(#\)](#)). Therefore, we incorporated these recent developments and modeled how the addition of dsRNA could disrupt the steady-state RNA metabolism of the targeted gene using ordinary differential equations (Fig. 4B [↗](#)). While there are many parameters that one could include in any model, we have used a conservative set of parameters for looking at the overall RNAi process without explicitly modeling sub-steps in detail. For example, production of 22G RNA is modeled as a single step rather than one that incorporates how the frequency of Cs in template mRNA, the subcellular localization of mRNA, secondary structure formation in mRNAs, etc. impact the efficiency of silencing. Similarly, genome sequence and its effect on transcription and/or splicing are modeled as a single step, rather than one that looks a frequency of repeats, sizes of introns, chromatin environment, etc. We expect that future studies will build upon this initial model hand-in-hand with the more sophisticated experimental tests needed to test such detailed hypotheses.

In this initial model (Fig. 4B [↗](#)), the steady-state levels of pre-mRNA and mRNA – which depend on production, maturation, and turnover – could be altered upon the addition of matching dsRNA through the generation of new RNA species (1° siRNA, 2° siRNA, pUG RNA) that are also subject to turnover. To accommodate these known intermediates and interactions, we used six differential equations to describe the rate of change of key RNA species (dsRNA (ds), 1° siRNA (pri), pUG RNA (ug), 2° siRNA (sec), pre-mRNA (p), and mRNA (m)) with rate or binding constants for different processes ($k1$ through $k9$), turnover rates for different RNAs (T_{pri} , T_{ug} , T_{sec} , T_p , T_m), and variables for the lengths of RNAs (l_{ds} – dsRNA; l_m – mRNA). For example, the rate of change over time for 1° siRNA is modeled as

$$k1.[ds].\frac{l_{ds}}{22} - k2.[pri].[m] - T_{pri}.[pri]$$

which includes the idea that 1° siRNAs are cleaved into 22nt sequences at a certain rate dependent on the dsRNA concentration ($k_{1,[ds],\frac{1}{22}}$), the amount of 1° siRNAs that can bind target transcripts (k_2 , $[pri]$, $[m]$), and the turnover of the 1° siRNAs (T_{pri} , $[pri]$), which notably is dependent on the concentration of the 1° siRNAs.

To illustrate the relative dynamics of different RNA species upon the addition of dsRNA, we computed the concentrations of dsRNA, 1° siRNA, pUG RNA, 2° siRNA, pre-mRNA, and mRNA using the equations after assigning arbitrary values for the different constants (**Fig. 4B** [↗](#); see legend for parameter values). To account for the non-negative values of all RNA species within cells, we ensured that the values of incremental change dx for any species x was only added if $(x+dx) > 0$ and set to be 0 if $(x+dx) \leq 0$. This bounding of the rate equations allows for any approach to zero. As expected, the levels of dsRNA decrease (**Fig. 4B** [↗](#), red) as it is processed into 1° siRNA (**Fig. 4B** [↗](#), purple), which eventually decays because of turnover. This transient accumulation of 1° siRNA is followed by that of pUG RNAs (**Fig. 4B** [↗](#), green) and of 2° siRNA (**Fig. 4B** [↗](#), brown). Silencing of the target is reflected in the lowered levels of mRNA (**Fig. 4B** [↗](#), blue) and pre-mRNA (**Fig. 4B** [↗](#), orange). However, these levels eventually recover upon turnover of the silencing intermediates (1° siRNA, pUG RNA, 2° siRNA). Although we assumed the turnover of 1° siRNA, 2° siRNA, and pUG RNA for the modeling, the experimental demonstration of recovery (either of individual RNA species or of the entire phenotype) from RNA silencing in non-dividing cells would be needed to support the existence of such turnover mechanisms for these different types of RNAs.

For any gene, the time to knockdown (t_{kd}) and the duration of knockdown (t_{kd}) could be used to evaluate the efficiency of RNAi (knockdown = 10% of initial mRNA concentration in **Fig. 4B** [↗](#)). The different RNA species made downstream of 1° RNA binding in *C. elegans* provide the opportunity for multiple parameters to differ between genes. Therefore, we varied each parameter and examined t_{kd} and t_{kd} as indicators of efficiency (Fig. S5). Overall, t_{kd} and t_{kd} were uncorrelated (**Fig. 4C** [↗](#)), with cases of rapid but transient knockdown, which would necessitate multiple dosing of dsRNA for sustained effects. While loss of function through the reduction of mRNA levels is often the intended goal of knockdown, RNA intermediates could serve as convenient and quantitative measures of molecular changes. For example, the abundant 2° siRNAs have been a widely used molecular indicator of silencing (e.g., (Gu et al., 2009 [↗](#))). However, the maximal amount of 2° siRNAs that accumulate is not correlated with strong silencing as measured by the minimal amount of mRNA during knockdown (**Fig. 4D** [↗](#)). Additionally, an increase in transcription generally resulted in poorer knockdown through changes in both t_{kd} and t_{kd} (**Fig. 4E** [↗](#)), consistent with the obvious expectation that a gene with transcriptional upregulation during exposure to dsRNA will be more difficult to knockdown.

Efficient silencing using dsRNA is possible in many organisms, including mammals, despite silencing relying on mostly post-transcriptional degradation of mRNA without the production of pUG RNA or 2° siRNA (Sandy et al., 2005 [↗](#)). To explore differences between genes that could impact the efficiency of RNA silencing universally in any system, we simulated knockdown through the post-transcriptional loss of mRNA alone by eliminating production of pUG RNAs, and thus downstream secondary small RNAs and transcriptional silencing (**Fig. 4B** [↗](#), $k_3 = 0$). When a fixed amount of dsRNA was exposed to different genes with the same amount of mRNA at steady state, genes with higher mRNA turnover rates showed less efficient knockdown (**Fig. 4F** [↗](#)). This inverse relationship is expected because to maintain the same steady state levels, genes with higher mRNA turnover must also have higher mRNA production. As a result, for the same amount of added dsRNA and the same steady-state level of mRNA before exposure to dsRNA, the mRNA levels will recover faster for genes with higher production coupled with higher turnover.

In summary, varying a few gene-specific parameters clarified the diversity of outcomes that are possible in response to the same dose of dsRNA. Gene-specific differences make the time to knockdown and the duration of knockdown uncorrelated and reduce the utility of key intermediates of RNA silencing as predictors of knockdown efficiency. Increases in transcription

during exposure to dsRNA and high turnover of mRNA coupled with high production at steady state reduce the efficiency of knockdown. While the predictions of the model include quantitative relationships that will require advances in the quantitative measurement of many steps during RNAi, the model also makes some qualitative predictions that can be immediately tested.

Changing *cis*-regulatory elements of a gene impacts its requirements for silencing by dsRNA

A key realization from the exploration of the dynamic model for RNAi is that pre-existing RNA regulation of a gene impacts the response to dsRNA (Fig. 4). However, the individual impacts of the many features of a gene that together set its RNA metabolism (e.g., promoter, 3' *cis*-regulatory regions, introns, genomic location, etc.) are usually unknown. Nevertheless, as tests of this possibility, we altered target genes using Cas9-mediated genome editing and examined changes, if any, in the genetic requirement for NRDE-3 for silencing by ingested dsRNA.

First, we swapped the 3' *cis*-regulatory regions of *bli-1* and *unc-22* (Fig. 5A, S6A). Animals with the *unc-22* 3' *cis* sequence in place of the *bli-1* 3' *cis* (*bli-1p::bli-1::unc-22* 3' *cis*) showed a much reduced dependence on NRDE-3 (Fig. 5B). Animals with the *bli-1* 3' *cis* sequence downstream of the coding sequence of *unc-22* (*unc-22p::unc-22::bli-1* 3' *cis*) showed substantial twitching even without *unc-22* RNAi (Fig. S6B). Yet, we were able to discern an enhancement upon addition of *unc-22* dsRNA. This enhancement was absent in animals lacking NRDE-3. Together, these results provide evidence for prior regulation (potentially mediated via the 3' UTR) impacting the genetic requirements for silencing.

Next, we used the *bli-1* 5' *cis*-regulatory regions (promoter) and the cyclin E 3' *cis*-regulatory regions to drive *gfp* expression in hypodermal cells (Fig. 5A; *bli-1p::gfp::cye-1* 3' *cis*). Animals with this gene showed expression of GFP in the hypodermis, which is most easily visible in the head region and is detectably silenced by ingested *gfp*-dsRNA in a wild-type background (Fig. 5C). Similar silencing was detectable in animals lacking NRDE-3 (Fig. 5D), but the silencing in animals lacking MUT-16 or RDE-10 was much weaker (Fig. 5D). These observations are in contrast to the observed lack of detectable silencing of the wild-type *bli-1* gene in response to *bli-1*-dsRNA in animals lacking NRDE-3, MUT-16, or RDE-10 (Fig. 1D and 2B), suggesting that the *bli-1* promoter is not sufficient to confer these requirements on all genes. Together, these results reveal that two different genes expressed under the same promoter within the same tissue can have different requirements for silencing in response to dsRNA (*bli-1* vs *gfp* under the control of *bli-1p*).

These initial attempts to change the requirements for the response to dsRNA by altering the pre-existing regulation of target genes encourage the exploration of additional factors predicted to differentially influence RNA silencing of different genes (Fig. 4, Fig. S5).

Gene expression can recover after knockdown despite the presence of amplification mechanisms

The dynamic model (Fig. 4) assumes that all key RNA intermediates (1° siRNA, 2° siRNA, and pUG RNA) are subject to turnover. If this assumption is true, animals should be able to recover from RNA silencing in non-dividing cells despite the production of abundant 2° siRNAs using RNA-dependent RNA polymerases. Experimental detection of the re-establishment of wild-type phenotype after a pulse of RNAi would provide evidence not only for the recovery of mRNA levels but also the subsequent production of functional protein. To test this possibility, we exposed wild-type animals to a 1-hr pulse of dsRNA matching the sensitive target *unc-22* and examined them for the Unc-22 defect every 24 hours (Fig. 6A). With this limited exposure to dsRNA, we observed only ~80% silencing after the first 24 hours, which reached ~100% by day 3, suggesting that it takes a couple of days after exposure to small amounts of dsRNA to observe complete silencing.




Fig. 5.

Changing *cis*-regulatory regions or open reading frames can alter the observed dependence on NRDE-3 for silencing by dsRNA.

(A) Schematics (as in Fig. 1) of the hybrid genes created to test the role of 3' and 5' *cis*-regulatory sequences in the response to dsRNA (see Methods for details). (B) The need for NRDE-3 to cause silencing using *bli-1*-dsRNA is reduced when the 3' *cis*-regulatory sequences of *bli-1* are replaced with that of *unc-22* (*bli-1(jam307)*). Fractions silenced, numbers scored, comparisons, asterisks, and error bars are as in Fig. 1 (C) Representative images (DIC, top; GFP fluorescence, bottom) of the head region of *bli-1p::gfp::cye-1 3'cis* animals either in response to a control vector (pL4440, left) or in response to *gfp*-dsRNA (right). The presence or absence of *gfp* in the head region was used to score the fraction silenced in (D). Scale bar = 100µm. (D) Use of the *bli-1* promoter alone does not confer the genetic requirements for silencing *bli-1*. Silencing of GFP expressed from the *bli-1p::gfp::cye-1 3'cis* transgene (*bli-1p::gfp(jam297)*) by *gfp*-dsRNA was measured in a wild-type background, in animals lacking NRDE-3 (*nrde-3(jam205)*), in animals lacking RDE-10 (*rde-10(jam206)*), in animals lacking MUT-16 (*mut-16(jam148)*), and in animals lacking both MUT-16 and RDE-10 (*mut-16(jam148) rde-10(jam196)*). Fractions silenced, numbers scored, comparisons, asterisks, and error bars are as in Fig. 1.

This delay could be driven by the time required for the buildup of RNA intermediates required for silencing (1° siRNA, 2° siRNA, and/or pUG RNA), for the turnover of UNC-22 protein, and/or for the dissipation of events downstream of the molecular role of UNC-22. Consistent with recovery, silencing was only observed in ~50% of the animals on day 5, which dropped to ~36% by the eighth day after RNAi. In contrast, animals that were continually fed *unc-22* RNAi showed ~100% silencing even at day 7 (**Fig. 6A**), suggesting that the RNAi machinery remains functional in aging animals. Thus, these results support the turnover of all key RNA intermediates generated during RNAi – 1° siRNA, 2° siRNA, and pUG RNA, and highlights for the first time that a target gene can recover from RNAi even in non-dividing cells despite their use of RNA-dependent RNA polymerases to amplify silencing signals.

Sequences in mRNA that match the trigger dsRNA are hot zones of pUG RNA production

Of the RNA intermediates generated during RNAi, pUG RNAs have been proposed to be used as stable templates to produce small RNAs (Shukla et al., 2020). Sustained production of small RNAs could occur if the targeting of mRNA by 2° siRNA resulted in further pUG RNA production, subsequent 3° siRNA production, and so on, thereby providing a way for silencing to persist despite the turnover of all RNA species. However, the production of such 3° siRNA has been observed only when targeting a germline gene (Sapetschnig et al., 2015) and not when targeting a somatic gene (Pak et al., 2012). To examine whether such repeated rounds of pUG RNA production occur during RNAi of *unc-22*, we fed wild-type worms bacteria that express *unc-22* dsRNA or control dsRNA (L4440) and looked for the presence of pUG RNAs. These RNAs are detected as a heterogenous mixture using RT-PCR with a poly-CA 3' primer and gene-specific 5' primers. Consistent with the production of pUG RNAs upon targeting by 1° siRNAs, we detected pUG RNAs generated after cleavage within the *unc-22* mRNA sequence that matches the dsRNA (**Fig. 6B**, 0kb 5' primer). Since 2° siRNAs are made with a 5' bias on the mRNA template (Pak et al., 2012; Pak and Fire, 2007), pUG RNAs generated in response to targeting by 2° siRNAs are expected to include mRNAs cleaved upstream of the sequence matching the dsRNA. Surprisingly, all pUG RNAs detected using a 5' primer ~1kb upstream of the target sequence were larger than 1 kb (**Fig. 6C**, 1kb 5' primer), suggesting that there is a dearth of pUG RNA formation through cleavage within 1 kb upstream of sequences targeted by dsRNA. Notably, this absence is despite the expected relative ease of amplifying shorter sequences when compared with amplifying longer sequences using the same primers. This lack of detectable pUG RNAs upstream suggests that, during RNAi in somatic cells, the addition of pUG tails is enriched within a zone on target mRNAs that share homology with the dsRNA trigger (“pUG zone”). This restricted production of pUG RNAs supports the idea that amplification is not perpetual and that mRNA levels can thus recover over time.

Discussion

Our results suggest that an intersecting network of regulators formed by the intrinsically disordered protein MUT-16, the Maelstrom-domain protein RDE-10, the nuclear Argonaute NRDE-3, and other Argonaute proteins can explain silencing of somatic targets by RNA interference despite stark differences in the genetic requirements for silencing different genes. The requirement for NRDE-3 can be overcome by enhanced dsRNA processing or by changing the cis-regulatory sequences of the target gene. However, the combined loss of both MUT-16 and RDE-10 eliminates all detectable silencing and this requirement cannot be overcome by enhancing dsRNA processing. Animals can recover from silencing in non-dividing cells, which supports the turnover of all key RNA intermediates (1° siRNA, 2° siRNA, and pUG RNA). Consistent with the ability to




Fig. 6.

Animals recover from a pulse of RNAi and production of pUG RNAs is restricted despite continuous exposure to dsRNA.

(A) Response to a pulse of feeding RNAi. (*top*) Schematic of assay. Animals were exposed to *unc-22* RNAi for one hour and then returned to OP50 plates. (*bottom*) A separate cohort of animals was scored for silencing after each subsequent 24hr period. Fractions silenced, numbers scored, comparisons, asterisks, and error bars are as in [Fig. 1](#). A weak Unc-22 defect indicates animals that were nearly completely still except for a slight twitch of the head or of the tail. (B) pUG RNA production in response to continuous exposure to *unc-22* dsRNA. (*top*) Schematic depicting the PCR primers used for the RT-PCR to detect pUG RNAs. Two sets of primers (0kb, purple; 1kb, blue) positioned 5' of the *unc-22*-dsRNA (orange), combined with an RT primer that contains nine CA repeats were used. (*bottom*) Populations of small RNAs that might be detected by pUG-PCR as above. Population I would suggest that amplified 2° siRNAs in addition to 1° siRNAs are capable of guiding target mRNA cleavage and poly-UG addition. Population II would suggest only 1° siRNAs can initiate pUG RNA production. (C) (*left*) Distribution of DNA amplified from pUG RNAs. Lanes with PCR products amplified from total RNA of animals fed *unc-22* dsRNA (*unc-22*) or L4440 dsRNA (control) isolated from three biological replicates each (1-3), or a no-template control (NTC) with no added RNA (-) are shown. Different bands are detected for each primer set (0kb, top vs. 1kb, bottom). A gene with a poly-UG sequence encoded in the genome (*gsa-1*) serves as a loading control. (*right*) Schematic of siRNA production in somatic cells. Successive rounds of amplified small RNAs would map progressively closer to the 5' end of the target transcript. Since we did not detect poly-UG addition upstream of the region homologous to the dsRNA trigger, it is unlikely that 3° siRNAs are being produced.

recover from silencing, unlimited rounds of siRNA amplification is curbed by restricting the cleavage and tailing of mRNAs for making pUG RNAs to ‘pUG zones’ that match the dsRNA sequence (see **Fig. 7** for an overview of findings).

Universal and gene-specific requirements for RNAi

RNAi requires the entry of dsRNA into cells, the processing of dsRNA into small RNAs, recognition of target mRNA, generation of additional small RNAs, and downstream gene silencing mechanisms. The upstream processes of entry, processing, and recognition do not depend on the mRNA being targeted and are thus presumably universal. Consistently, the dsRNA importer SID-1, the endonuclease DCR-1, and the primary Argonaute RDE-1 are required for all RNAi. In contrast, since the mRNA is used as a template to generate the abundant secondary small RNAs in *C. elegans* (Pak et al., 2012) or additional dsRNAs in other systems (e.g., in plants (Sanan-Mishra et al., 2021)), the silencing of different mRNAs could diverge through the selective recruitment of different collections of regulators. In support of this possibility, the two model genes we analyze in this study, *unc-22* and *bli-1*, show stark differences in the requirements of some RNAi factors for silencing (**Fig. 1**). While these differences could be attributed to their expression in different tissues, the ability to bypass some requirements (**Fig. 3**) argues against this possibility. Specifically, if the requirement for NRDE-3 for silencing *bli-1* (hypodermal gene) but not *unc-22* (muscle gene) is because of the lack of a parallel regulator in the hypodermis but not in the muscle, then enhancing dsRNA processing would be unable to bypass the NRDE-3 requirement (**Fig. 3**). Neither the silencing of *dpy-7* nor the silencing of *bli-1p::gfp* requires NRDE-3 for silencing despite their expression in the hypodermis (**Fig. 2F**, **Fig. 5D**). Strikingly, changing the 3' cis-regulatory sequences of *bli-1* also made its silencing largely independent of NRDE-3 (**Fig. 5B**), providing direct evidence for the prior regulation of a gene dictating the genetic requirements for silencing in response to dsRNA. The fact that any two of MUT-16, NRDE-3, and RDE-10 – three very structurally and functionally different proteins – are required for *unc-22* silencing suggests that each of these proteins could be contributing to silencing of any RNAi target. Despite this potential use of an intersecting network for silencing all somatic genes, different genes could critically depend on different regulators because of differences in their mRNA metabolism and/or subcellular localization (summarized as threshold differences in **Fig. 2**). Intermediate steps that require the participation of mRNA such as the production of 2° siRNA could have complex dependencies, making RNA intermediates poor predictors of silencing efficiency (**Fig. 4D**). For example, the subcellular localization of mRNA could increase or decrease its interaction with RdRPs and thus influence the levels of 2° siRNAs made. Future studies that address the dynamics and subcellular localization of target mRNA before RNAi and the subcellular localization of components of the RNAi machinery are required to test these hypotheses.

Production of 2° siRNAs

Multiple small RNA species of defined lengths and 5'-nt bias have been detected in *C. elegans*. Of these, 22G RNAs (2° siRNAs) are the most abundant and arise from amplification downstream of exposure to dsRNA and in multiple endogenous small RNA pathways (Gu et al., 2009). Our results suggest that production of 2° siRNAs in response to dsRNA is eliminated in animals that lack both MUT-16 and RDE-10 (**Fig. 3C**). While the precise mechanisms of amplification are unknown, MUT-16 is thought to nucleate perinuclear foci in the germline (Phillips et al., 2012; Uebel et al., 2018) that recruit many additional components – RDE-2, MUT-7, MUT-14, MUT-15, NYN-1, NYN-2, RDE-8, RDE-3/MUT-2 etc. (Phillips and Updike, 2022; Uebel et al., 2018). Similar interactions may occur in somatic cells despite the lack of detectable perinuclear foci. The roles of most proteins remain obscure, but RDE-8 (Tsai et al., 2015) and RDE-3/MUT-2 (Shukla et al., 2020) have demonstrated roles in the cleavage and pUGylation of mRNAs, respectively. Yet, the observation of silencing in the absence of MUT-16 that is eliminated upon additional loss of RDE-10 suggests that RDE-10 and associated proteins (e.g., RDE-11, RDE-12) play an independent role




Figure 7.

Overview of findings.

(left) Amplification of small RNAs occurs through an intersecting network of regulators (orange arrows) and in response to somatic RNAi, the addition of poly-UG repeats (green) is restricted to a 'pUG zone' (red) that is homologous to the dsRNA trigger. (right) In response to a pulse of dsRNA (red), levels of the target mRNA (blue) recover, suggesting that the RNA silencing intermediates (1° siRNAs, purple; pUG RNAs, green; 2° siRNAs, brown) undergo turnover.

in the amplification of 2° siRNAs (Yang et al., 2012 [↗](#); Zhang et al., 2012 [↗](#)). The subcellular localization of RDE-10 and whether small RNAs that require RDE-10 for production also rely on RDE-8 and RDE-3/MUT-2 as expected for amplification using pUG RNA templates remains to be determined.

Loss of RDE-10 reduces the production of 22G RNAs downstream of exogenous dsRNA and downstream of endogenous small RNAs called 26G RNAs that are 26-nt long and have a 5' G (Yang et al., 2012 [↗](#); Zhang et al., 2012 [↗](#)). Current models for the production of 26G RNAs (Blumenfeld and Jose, 2015 [↗](#); Chaves et al., 2021 [↗](#)) propose that the RdRP RRF-3 transcribes long antisense RNA from internal C nucleotides on template mRNA, the phosphatase PIR-1 converts the 5' triphosphate of the RdRP product into 5' mono phosphate, the template is then trimmed by the 3'-5' exonuclease ERI-1 to generate a blunt-ended dsRNA, which is then cleaved by DCR-1 to generate the mature 26G RNAs that are bound by the Argonaute ERGO-1. While a similar preference by RdRPs can explain the 5'G bias of the downstream 22G RNAs, the mechanism(s) for generating RNA that are precisely 22 nucleotides long remain unclear. This precision could be achieved either through the trimming of template mRNAs into 22-nt long pieces or through the trimming of secondary small RNAs made by RdRPs into 22-nt long pieces. The detection of long pUG RNAs with no detectable shorter pUG RNAs upstream of sequences matching the dsRNA (Fig. 6C [↗](#)) argues against the 3' trimming of mRNA templates to generate shorter RNAs that then get pUGylated to become stabilized templates for RdRPs and against pUG RNA generation driven by successive rounds of 22G RNA production in somatic cells. Furthermore, potential 5' trimming or endonucleolytic cleavage of long pUG RNA to generate a 22-nt template for RdRPs cannot explain the 5'G bias of 22G RNAs. Since Argonautes bind the 5' end of small RNAs and can associate with RNAs of different lengths (Ruby et al., 2006 [↗](#)), we suggest a model whereby RDE-10 and downstream Argonautes together play a role in the maturation of 22-nt siRNAs from longer RdRP products.

RDE-10 has a conserved Maelstrom domain that shares homology with the DnaQ-H 3'-5' exonuclease family (Zhang et al., 2008 [↗](#)) and the mutation we identified as disrupting silencing by dsRNA (Fig. 1D [↗](#)) alters a residue located near the highly conserved ECHC zinc-binding motif (Fig. S1C). Intriguingly, the Maelstrom domain of RDE-10 shares high structural homology with the 3'-5' exonuclease domain of ERI-1 (Fig. S7C, *left*) but not the exonuclease domain of MUT-7 (Fig. S7C, *right*). ERI-1 can trim single-stranded RNA overhangs in vitro (25), and is required for the production of 26G RNAs (Duchaine et al., 2006 [↗](#)) and for the maturation of rRNAs (Gabel and Ruvkun, 2008 [↗](#)). While no 3'-5' exonuclease activity of RDE-10 or its orthologs has been demonstrated, Maelstrom domain-containing proteins in insects exhibit single-stranded RNA endonuclease activity in vitro (Matsumoto et al., 2015 [↗](#)). Furthermore, RDE-10 could interact with other parts of the RNA silencing machinery (e.g., the Argonaute ERGO-1 as seen using immunoprecipitation (Yang et al., 2012 [↗](#); Zhang et al., 2012 [↗](#))) to recruit nucleases (e.g., NYN family exonucleases such as ERI-9 (Tsai et al., 2015 [↗](#))) that trim pre-22G RNAs to the 22-nt size preferred by Argonaute proteins. In support of such exonucleolytic trimming in conjunction with Argonaute binding, the 3'-5' exonuclease SND1 has been shown to trim the 3' ends of miRNAs bound to AGO1 in *Arabidopsis* (Chen et al., 2018 [↗](#)). Furthermore, piRNA maturation in *Drosophila* and mice suggests a model where piwi-type Argonautes bind the 5' end of the pre-piRNA followed by endonucleolytic cutting and exonucleolytic trimming to generate consistently sized mature piRNAs (Stoyko et al., 2022 [↗](#)). Finally, human ERI1 can trim Ago2-bound micro RNAs to 19-nt (Sim et al., 2022 [↗](#)).

Therefore, we propose that the production of 22G RNAs in response to the addition of dsRNA occurs as follows: (1) non-processive RdRPs (e.g., RRF-1 (Aoki et al., 2007 [↗](#))) make a heterogeneous mixture of short RNAs, (2) 2° Argonautes bind the 5' end of these pre-secondary siRNA, (3) RDE-10 and/or associated protein(s) remove excess 3' sequence to generate 22-nt siRNAs that are

effectively retained by the mature siRNA-Argonaute complex. Similar mechanisms could be used to generate other 22G RNAs that are independent of RDE-10 (Yang et al., 2012 [↗](#); Zhang et al., 2012 [↗](#)). Future studies are needed to test each aspect of the model.

Trade-offs in RNA interference

RNAi is now a widely applied tool for gene silencing in plants, insects, and humans. Like *C. elegans*, plants (Sanan-Mishra et al., 2021 [↗](#)) and some insects (Pinzón et al., 2019 [↗](#)) have RdRPs that could be used to make 2° siRNAs, but many other animals, including humans, do not have RdRPs and thus are unlikely to produce 2° siRNAs. However, silencing can fade despite the production of 2° siRNAs (Fig. 6A [↗](#)), highlighting the importance of dosage for all systems. Two parameters of importance for the acute efficacy of any dsRNA-based drug are the time to knockdown (t_{kd} in Fig. 4B [↗](#)) and duration of knockdown (t_{kd} in Fig. 4B [↗](#)). The various values of t_{kd} that are possible for each t_{kd} (Fig. 4C [↗](#)) cautions against using a rapid onset of silencing (low t_{kd}) as the sole indicator of promise during early stages of drug development when long-term effects of a drug are often not evaluated in the interest of expedience. In short, a drug that takes longer to cause an effect could have a more long-lasting effect. Since a dsRNA drug can be synthesized for any target with equal effort, considerations for the choice of target could be worthwhile because differences in RNA metabolism between two targets of equal importance can influence the efficacy of the dsRNA drug in all systems. If two genes are at steady state, then the gene with higher mRNA turnover will be more difficult to knockdown because of higher rates of mRNA production (Fig. 4F [↗](#)). Similarly, in the absence of a steady state, a gene undergoing upregulation of transcription, splicing, and/or mRNA export during the administration of the drug will be difficult to knockdown (e.g., Fig. 4E [↗](#)).

In the longer term, a concern for any drug is the development of resistance. When a gene with a high threshold for silencing is targeted, it could rely on multiple regulators that act in parallel to contribute to silencing (e.g., *bli-1* in this study), making resistance through the mutation of any one regulator more likely and necessitating another round of drug development. In contrast, genes with a lower threshold may not require all the regulators for silencing (e.g., *unc-22* in this study), making them ideal targets that remain silenced despite single mutations in many regulators of RNAi (e.g., RDE-10, MUT-16, or NRDE-3 in this study). These trade-offs inform the choice of therapeutic targets and dosage for sustained use of dsRNA-based drugs in agriculture and in human health. Anticipating mechanisms for the development of resistance before widespread use of an RNAi-based drug or treatment will be crucial for avoiding futile cycles of innovation. The ideal drug would require a minimal dose and use multiple intersecting paths to silence the target gene.

Materials and methods

Summary

All strains (Table S1) were grown at 20°C on Nematode Growth Medium (NGM) plates seeded with OP50 *E. coli* (Brenner, 1974 [↗](#)). Strains with mutations were generated through a genetic screen after mutagenesis using N-ethyl-N-nitrosourea (ENU), using standard genetic crosses (Brenner, 1974 [↗](#)), or using Cas9-mediated genome editing (Arribere et al., 2014 [↗](#); Dokshin et al., 2018 [↗](#); Paix et al., 2015 [↗](#)). Mutations induced upon ENU exposure were identified using whole genome sequencing (Illumina) followed by analyses of the resultant fastq files. Simulations of the RNAi response were used to identify the domain and range of values consistent with experimental data (Fig. 2 [↗](#)) and to explore parameters that support silencing (equilibrium model (Fig. S4) and dynamic model (Figs. 4 [↗](#) and S5)). Feeding RNAi experiments were performed by exposing worms to bacteria that express dsRNA (Kamath et al., 2003 [↗](#); Timmons and Fire, 1998 [↗](#)) either continuously or for a brief period (Fig. 6A [↗](#)). Multiple sequence alignment (Fig. S7) was performed using Clustal Omega (Sievers et al., 2011 [↗](#)) and manually annotated using Illustrator (Adobe).

Comparisons of protein structures were performed using AlphaFold predictions (Jumper et al., 2021 [↗](#); Varadi et al., 2022 [↗](#)), pair-wise alignment on Protein Data Bank (Zhang, 2005 [↗](#)), and the PyMOL Molecular Graphics System (v. 2.4.1 Schrödinger, LLC). Levels of *rde-4* mRNA (Fig. S3) and pUG RNA (Fig. 6C [↗](#)) were measured using reverse-transcription followed by polymerase chain reaction (RT-PCR). Transgenic strains that express *rde-1(+)* and *rde-4(+)* in specific tissues were generated using Mos1-mediated single copy insertion (MosSCI, (Frøkjær-Jensen et al., 2012 [↗](#)). Oligonucleotides used are in Table S2. Exact p-values and additional details for each experiment are in Table S3. All code used (R, Python, and Shell) is available at https://github.com/AntonyJose-Lab/Knudsen_et_al_2023 [↗](#).

Strains and oligonucleotides used

All strains (listed in Table S1) were cultured on Nematode Growth Medium (NGM) plates seeded with 100 µl of OP50 *E. coli* at 20°C and strains made through mating were generated using standard methods (Brenner, 1974 [↗](#)). Oligonucleotides used are in Table S2. Strains generated using Mos1-mediated Single Copy Insertion (MosSCI, (Frøkjær-Jensen et al., 2012 [↗](#))) of *rde-4* or *rde-1* rescues in the germline (as in (Marré et al., 2016 [↗](#))) or of *rde-4* rescues in the hypodermis (Raman et al., 2017 [↗](#)) were used in this study.

Genetic screen

This screen was performed by mutagenizing a strain (AMJ174) with the transgene *T(oxSi487[mex-5p::mCherry::H2B::tbb-2 3'UTR::gpd-2 operon::GFP::H2B::cye-1 3'UTR + unc-119(+)]*, (Devanapally et al., 2021 [↗](#))) silenced for >200 generations after introducing a mutation in *lin-2(jam30)* (sgRNA (P1), primers (P2, P3, P4) using Cas9-mediated genome editing of AMJ844 (*iT; dpy-2(e8)*, (Devanapally et al., 2021 [↗](#))) while correcting the *dpy-2(e8)* mutation to wild type (creating *dpy-2(jam29)*; sgRNA (P5), primers (P6, P7, P8)). The *lin-2* mutation limits brood size (Ferguson and Horvitz, 1985 [↗](#)) and facilitates screening. Near-starved animals (P0) of all life stages were mutagenized using 1mM N-Ethyl-N-Nitrosourea (ENU, Toronto Research Chemicals) for 4-6 hours. Mutagenized animals were washed four times with wash buffer (0.01% Triton X-100 in M9) and 2-3 adult animals were placed on NG plates seeded with OP50. Over the next 3 weeks, F1, F2, and F3 progeny were screened to isolate mutants that show mCherry fluorescence. These animals were singled out (up to 7 animals from each P0 plate) and tested for the persistence of expression in descendants. Of the 15 fertile mutants isolated using this primary screen, five with mutations in *mut-16* were analyzed in this study.

Whole genome sequencing

Libraries were prepared using TruSeq DNA Library Prep kits (Illumina) and samples were sequenced at Omega Biosciences. The fastq files obtained after Illumina sequencing (1x PE 150 b, Omega Biosciences) were analyzed to identify candidate mutations responsible for the observed defects in the sequenced strains. For each strain, sequences were trimmed using cutadapt (v. 3.5), mapped to the *C. elegans* genome (WBcel235/ce11) using bowtie2 (v. 2.4.2), sorted using samtools (v. 1.11), and the resulting.bam file was analyzed to call variants using snpEff (v. 5.0e). The variants classified as 'HIGH' or 'MODERATE' in the .ann.vcf file for each strain that were not shared by any two or more strains were culled as new mutations caused by mutagenesis in each strain. These new mutations in each strain were compared with those of all other strains ('in silico complementation') using a custom script to identify sets of strains with different mutations in the same genes. Specific details for each step are provided within the scripts '1_fastq_to_sorted_bam.sh', '2_sorted_bam_to_mutated_genes.sh', '3_in_silico_complementation.sh' available at GitHub (https://github.com/AntonyJose-Lab/Knudsen_et_al_2023 [↗](#)). Raw fastq files for the strains analyzed in this study (AMJ1023, AMJ1025, AMJ1035, AMJ1042, and AMJ1091) have been submitted to SRA (PRJNA928750).

Modeling and simulation

The RNAi response was explored using three models of increasing complexity: (1) a single-network model of protein factors with branching pathways for RNA amplification and subsequent gene silencing (Fig. 2); (2) an equilibrium model for the dependence of mRNA and pre-mRNA on small RNAs and other RNA intermediates (Fig. S4); and (3) a dynamic model using ordinary differential equations for the dependence of mRNA and pre-mRNA on small RNAs and other RNA intermediates (Fig. 4 and S5). Simulations of single network and exploration of equilibrium model were conducted in R (v. 3.6.3). Simulations of the dynamic model were conducted in Python (v. 3.8.5) and in R (v. 4.1.0).

Intersecting network

Random numbers from 0 to 2 were selected for each of the assigned variables (Nm , Nr , Om , Or) and parameter sets that satisfy experimental constraints were plotted. Specific details are provided within the script '2022_6_13_RNAi_in_Celegans_linear_modified.R' available at GitHub (https://github.com/AntonyJose-Lab/Knudsen_et_al_2023).

Equilibrium model

This model for RNAi interference assumes that all reactions have reached equilibrium. Additional assumptions include (1) 1° siRNAs, then pUG RNAs, then 2° siRNAs are made sequentially, (2) no 3° siRNAs are produced for these somatic targets (supported by (Pak et al., 2012)), (3) there is no recycling of full-length mRNA or full-length pre-mRNA after small RNA binding, i.e., multiple rounds of binding by different small RNAs to the same intact mRNA or pre-mRNA molecules is not allowed and (4) there are no other mechanisms for the turnover of the RNA species considered in the timescale considered. Specific details are provided within the script '2022_2_9_RNAi_network_thresholds_simpler.R' available at GitHub (https://github.com/AntonyJose-Lab/Knudsen_et_al_2023).

Dynamic model

A series of differential equations were used to describe the rate of change for dsRNA, 1° siRNAs, mRNAs, pre-mRNAs, pUG RNAs, and 2° siRNAs, and numerically simulated using the 4th Order Runge-Kutta method. Specific details are provided within the scripts '2022_6_29_Celegans_RNAi_ODEs_RK4_method_d6.py' and '2022_7_14_RNAiDynamics_ODEs_Parameter_Analysis.R' available at GitHub (https://github.com/AntonyJose-Lab/Knudsen_et_al_2023).

Genome editing

The gonads of adult *C. elegans* were injected with nuclear-localized Cas9 (PNA Bio) preincubated at 37°C for 10 min with a hybridized crRNA/tracrRNA (Integrated DNA Technologies), as well as an oligonucleotide or PCR-amplified homology repair template. Plates with successfully edited F1 animals were identified by screening the Dpy or Rol animals obtained when using *dpy-10* editing as a co-CRISPR (Arribere et al., 2014; Paix et al., 2015) or for Rol animals when using the pRF4 plasmid as a co-injection marker (Dokshin et al., 2018).

To introduce a premature stop codon in *mut-16*

Injection of a crRNA with the target sequence (P9) (Integrated DNA Technologies), tracrRNA, Cas9, a *mut-16(-)* homology repair template (P10) mimicking the mutation in *mut-16(jam139)*, predicted amino acid change Y294*, and *dpy-10* crRNA (P11) and *dpy-10(-)* homology repair template (P12) into N2 or AMJ1489 and subsequent screening were performed as described above. Genotyping for *mut-16(jam148, jam240, jam265, jam266, jam267, or jam268)* was performed using duplex PCR

(P13, P14) followed by restriction digestion with BstBI. The nonsense mutations in different strains (AMJ1397, AMJ1611, AMJ1672, AMJ1673, AMJ1674, and AMJ1675) were verified by Sanger sequencing.

To make the *mut-16(-) rde-10(-)* double mutant

Injection of a crRNA with the target sequence (P15) (Integrated DNA Technologies), tracrRNA, Cas9, a *rde-10(-)* homology repair template (P16) mimicking the mutation in *rde-10(jam248)*, and *dpy-10* crRNA (P11) and *dpy-10(-)* homology repair template (P12) into AMJ1397 (*mut-16(jam148)*) and subsequent screening were performed as described above. Genotyping for *rde-10(-)* was performed using duplex PCR (P17, P18) followed by restriction digestion with EcoRV. A strain with a mutation in *rde-10* that results in a 115-bp frameshift followed by an early stop codon was designated as AMJ1470.

To introduce the mutation in *rde-10* that will encode RDE-10(Ser228Phe)

Injection of a crRNA with the target sequence (P15) (Integrated DNA Technologies), tracrRNA, Cas9, a *rde-10(-)* homology repair template (P16) mimicking the mutation in *rde-10(jam248)* (Fig. S1), and pRF4 into N2 and subsequent screening were performed as described above. Genotyping for the mutation was performed using duplex PCR (P17, P18) followed by restriction digestion with EcoRV. A strain with the missense mutation verified by Sanger sequencing was designated as AMJ1489.

To introduce a premature stop codon in *rde-10*

Injection of a crRNA with the target sequence listed as (P19) (Integrated DNA Technologies), tracrRNA, Cas9, a *rde-10(-)* homology repair template (P20) (predicted amino acid change Q73*) and pRF4 into N2 and subsequent screening were performed as described above, Genotyping for *rde-10(-)* was performed using duplex PCR (P21, P22) and restriction digestion with DpnII to isolate the mutant from N2. A strain with a 2-bp deletion near Q73 that results in a frameshift and an early stop codon was designated as AMJ1614.

To introduce a premature stop codon in *nrde-3*

Injection of a crRNA with the target sequence (P23) (Integrated DNA Technologies), tracrRNA, Cas9, a *nrde-3(-)* homology repair template (P24), mimicking *nrde-3(gg066)* (25), and pRF4 into N2 and subsequent screening were performed as described above. Genotyping for *nrde-3(jam205)* was performed using duplex PCR (P25, P26) followed by restriction digestion with AclI. A strain with the nonsense mutation verified by Sanger sequencing was designated as AMJ1510.

To introduce a premature stop codon in *eri-1*

Injection of a crRNA with the target sequence (P27) (Integrated DNA Technologies), tracrRNA, Cas9, an *eri-1(-)* homology repair template (P28), predicted to encode ERI-1(E225*) after the edit, and pRF4 into AMJ1611 or N2 and subsequent screening were performed as described above. Genotyping for *eri-1(jam260, jam261, jam262, jam263, or jam264)* was performed using duplex PCR (P29, P30) followed by restriction digestion with DpnII. Additionally, when *eri-1(mg366)* was crossed with other mutants, duplex PCR with P31 and P32 was used for genotyping.

To create a transgene with the *bli-1* promoter

Injection of two crRNAs with the target sequences (P33, P34) (Integrated DNA Technologies), tracrRNA, Cas9, a homology repair template that was amplified using sequences (P35, P36) (Phusion High-Fidelity DNA Polymerase, New England BioLabs), which amplifies the promoter region of *bli-1*, and pRF4 into AMJ1195 [*mex-5p::gfp::cye-1 3' utr*] (40) and subsequent screening were performed as described above. Genotyping for *bli-1p::gfp* was performed using triplex PCR (P37, P38, P39). Additional genotyping after crosses was done using triplex PCR with sequences

(P40, P41, P42). The resulting strain (AMJ1709) resulted in successful integration of ~75% of the *bli-1* promoter upstream of the *mex-5* promoter, and showed GFP expression within the hypodermis (most notable in the head region, see [Fig. 5C](#)) and in the germline (data not shown).

To mutate the 3' cis-regulatory regions of *bli-1*

Injection of two crRNAs with the target sequences (P43, P44) (Integrated DNA Technologies), tracrRNA, Cas9, a homology repair template that was amplified using sequences (P45, P46) (Phusion High-Fidelity DNA Polymerase, New England BioLabs), which amplifies the 3'utr + 50bp of *unc-22*, and pRF4 into N2 and subsequent screening were performed as described above. Genotyping for the altered *bli-1* gene was performed using triplex PCR (P47, P48, P49). A strain with partial (~65%) integration of the *unc-22* 3' cis region, a repeat of the first 183 bp of the *unc-22* 3' cis region, and the endogenous *bli-1* 3' cis region was designated as AMJ1754 and a strain with complete integration was designated as AMJ1755.

To mutate the 3' cis-regulatory regions of *unc-22*

Injection of two crRNAs with the target sequences (P50, P51) (Integrated DNA Technologies), tracrRNA, Cas9, a homology repair template that was amplified using sequences (P52, P53) (Phusion High-Fidelity DNA Polymerase, New England BioLabs), which amplifies the 3'utr + 50bp of *bli-1*, and pRF4 into N2 and subsequent screening were performed as described above. Genotyping for the altered *unc-22* gene was performed using triplex PCR (P47, P48, P54). Strains with mutated *unc-22* 3' cis region were isolated and designated as AMJ1730 and AMJ1731.

Sequence and structure alignments

Sequences of *C. elegans* proteins were obtained from WormBase; sequences of proteins from all other species were obtained from UniProt. Alignments were created using Clustal Omega (EMBL-EBI) with default settings.

PyMOL (v. 2.4.1) was used to modify and annotate PDB files. The RDE-10 (UniProt: Q9N3S2) PDB file is based on predictions from AlphaFold. Protein domains were colored based on homology to domains as found in the EMBL-EBI Pfam database (Maelstrom: PF13017). The protein structure alignment was done using the Pairwise Structure Alignment from Protein Data Bank with rigid parameters (RMSD Cutoff 3; AFP Distance Cutoff: 1600; Fragment length: 8). The exonuclease domain of ERI-1 (UniProt:O444606) and of MUT-7 (UniProt:P34607) were compared with the Maelstrom domain of RDE-10.

Feeding RNAi

Control RNAi by feeding *E. coli* containing the empty dsRNA-expression vector (L4440), which can generate a short dsRNA derived from the multiple cloning site but does not produce dsRNA against any *C. elegans* gene, was done in parallel with all RNAi assays.

P0 and F1 feeding

Bacteria expressing dsRNA was cultured in LB media with 100 µg/µl carbenicillin overnight at 37°C at 250 rpm. 100 µl of cultured bacteria was then seeded onto RNAi plates [NG agar plate supplemented with 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) (Omega) and 25 µg/ml carbenicillin (MP Biochemicals)]. Adult animals were passaged onto seeded RNAi plates and removed after 24hr. Progeny were scored for silencing by bacteria expressing dsRNA targeting *unc-22* (defect evident as twitching within ~3 min. in 3 mM levamisole) or *bli-1* (defect evident as blisters along the body).

P0 pulse feeding

L4 and young adult animals were placed on seeded RNAi plates for 1hr after which they were transferred to an OP50 plate for 1hr, and then transferred to a new OP50 plate once again to minimize the residual RNAi food carryover. Animals were left on OP50 plates and scored every 24hr for 8 subsequent days with transfer to new OP50 plates every two days to prevent overcrowding.

F1 only feeding

A single L4 or young adult (1 day older than L4) animal (P0) was placed on an RNAi plate seeded with 5µl of OP50 and allowed to lay eggs. After 1 day, when most of the OP50 was eaten, the P0 animal was removed, leaving the F1 progeny. 100µl of an overnight culture of RNAi food (*E. coli* which express dsRNA against a target gene) was added to the plate. Two or three days later, the F1 animals were scored for gene silencing by measuring gene-specific defects.

RNA extraction and PCR with reverse transcription (RT-PCR)

Total RNA was extracted using TRIzol (Fisher Scientific) from pellets of mixed-stage animals collected from non-starved but crowded plates in biological triplicate for each strain after exposure to either *unc-22* RNAi or the L4440 vector. The aqueous phase was then washed with an equal amount of chloroform and precipitated overnight at -20°C with 10 µg glycogen (Invitrogen) and 1 ml of isopropanol. RNA pellets were washed twice with 70% ethanol and resuspended in 25 µl of nuclease free water.

RT-PCRs for pUG RNAs (**Fig. 6C**) were done as described earlier (Shukla et al., 2020). Briefly, cDNA was created from isolated RNA using Superscript III Reverse Transcriptase (ThermoFisher) and a universal primer (P55) that contains nine CA repeats and two adapter sequences. The cDNA was used as a template for a Phusion (NEB) PCR with the first set of primers (adapter 1 P56; *gsa-1* P57; 0kb P58; 1kb P59). The amplicon was diluted 1:20 and used as template for the nested Phusion PCR with the second set of primers (adapter 2 P60; *gsa-1* P61; 0kb P62; 1kb P63). The final PCR product (20 µl) was loaded on a 1% agarose gel and imaged. An annealing temperature of 60°C was used for *gsa-1* and 57°C was used for all other primer sets.

For semi-quantitative RT-PCR (Fig. S3), RNA from each strain was isolated from 50 L4-staged animals as described earlier (Jobson et al., 2015). Primer P64 was used to reverse transcribe the sense strand of *rde-4* and P65 was used to reverse transcribe the sense strand of *tbb-2*. The resulting cDNA was used as a template for PCR (30 cycles for both *rde-4* and *tbb-2*) using Taq polymerase and gene-specific primers (P66, P67 for *rde-4* and P68, P69 for *tbb-2*). Intensities of the bands were quantified using Image J (NIH). The relative intensity of the *rde-4* band normalized to that of the *tbb-2* band was set as 1.0 in wild type. The relative normalized intensity of the *rde-4* band in WM49 (*rde-4(ne301)*) was subtracted from that in AMJ565 to report the levels of *rde-4(+)* mRNA (0.3 relative to wild type).

Microscopy

Following *gfp* RNAi (**Figs. 5C**, **5D**), F1 adult animals were mounted in 10 µl of 3 mM levamisole on a 2% agar pad and imaged under a coverslip using a Nikon AZ100 microscope and Prime BSI Express sCMOS camera. A C-HGFI Intensilight Hg Illuminator was used to excite GFP (filter cube: 450 to 490 nm excitation, 495 dichroic, and 500 to 550 nm emission). Representative images for GFP expression were adjusted to identical levels in Fiji (NIH) for presentation.

Rationale for inferences

Prior knowledge

Gene-specific requirements for RNA silencing could reflect specialization along pathways, as is supposed for multiple endogenous small RNA pathways in *C. elegans*. Reasons that impact the efficiency of silencing a gene are obscure because of a lack of a quantitative model for RNAi that incorporates recently discovered RNA intermediates.

Evidence supporting key conclusions

Three different proteins, MUT-16, RDE-10, and NRDE-3, play a role in RNAi such that each is singly required for silencing *bli-1* but any two are sufficient for silencing *unc-22*. Simulations support the parsimonious hypothesis that this difference in requirements can be explained by quantitative contributions from regulators within an intersecting network for silencing both genes but not by parallel pathways downstream of mRNA recognition. Consistently, the requirement for NRDE-3 for silencing *bli-1* is bypassed by enhancing the processing of dsRNA through the loss of ERI-1 or the overexpression of RDE-4.

A quantitative model for RNAi of any gene at steady state reveals several ways that differences in genetic requirements could arise for silencing different genes. Experimental tests that confirm predictions of the quantitative models include changes in the requirement for NRDE-3 for silencing caused by altering *cis*-regulatory regions of the gene targeted by dsRNA; recovery from silencing in non-dividing cells after exposure to a pulse of *unc-22* dsRNA, which supports the turnover of all key RNA intermediates (1° siRNAs, 2° siRNAs, and pUG RNAs) through mechanisms that are currently unknown; and the dearth of pUG RNA generation by 2° siRNAs, consistent with a lack of 3° siRNAs.

Data availability

All strains used are available upon request. Fastq files from whole-genome sequencing are available at NCBI SRA database with the accession number PRJNA928750.

Supplementary data

This manuscript includes supplementary data

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

We thank Mary Chey and Julianna Gross for some analysis of mutants from the genetic screen; Zhongchi Liu, Leslie Pick, Norma Andrews, and members of the Jose Lab for comments on the manuscript; and the *Caenorhabditis elegans* Genetic Stock Center for some worm strains.

Funding

This work was supported in part by National Institutes of Health Grants R01GM111457 and R01GM124356, and National Science Foundation Grant 2120895 to A.M.J.

References

1. Aoki K, Moriguchi H, Yoshioka T, Okawa K, Tabara H (2007) **In vitro analyses of the production and activity of secondary small interfering RNAs in *C. elegans*** *The EMBO Journal* **26**:5007–5019 <https://doi.org/10.1038/sj.emboj.7601910>
2. Arribere JA, Bell RT, Fu BXH, Artiles KL, Hartman PS, Fire AZ (2014) **Efficient Marker-Free Recovery of Custom Genetic Modifications with CRISPR/Cas9 in *Caenorhabditis elegans*** *Genetics* **198**:837–846 <https://doi.org/10.1534/genetics.114.169730>
3. Bergstrom CT, McKittrick E, Antia R (2003) **Mathematical models of RNA silencing: Unidirectional amplification limits accidental self-directed reactions** *Proceedings of the National Academy of Sciences* **100**:11511–11516 <https://doi.org/10.1073/pnas.1931639100>
4. Blumenfeld AL, Jose AM (2015) **Reproducible features of small RNAs in *C. elegans* reveal NU RNAs and provide insights into 22G RNAs and 26G RNAs** *RNA* <https://doi.org/10.1261/rna.054551.115>
5. Brenner S (1974) **THE GENETICS OF CAENORHABDITIS ELEGANS** *Genetics* **77**:71–94 <https://doi.org/10.1093/genetics/77.1.71>
6. Chaves DA, Dai H, Li L, Moresco JJ, Oh ME, Conte D, Yates JR, Mello CC, Gu W (2021) **The RNA phosphatase PIR-1 regulates endogenous small RNA pathways in *C. elegans*** *Molecular Cell* **81**:546–557 <https://doi.org/10.1016/j.molcel.2020.12.004>
7. Chen C-CG, Simard MJ, Tabara H, Brownell DR, McCollough JA, Mello CC (2005) **A Member of the Polymerase Beta Nucleotidyltransferase Superfamily Is Required for RNA Interference in *C. elegans*** *Current Biology*
8. Chen J *et al.* (2018) **Structural and biochemical insights into small RNA 3' end trimming by *Arabidopsis* SDN1** *Nat Commun* **9** <https://doi.org/10.1038/s41467-018-05942-7>
9. Das PR, Sherif SM (2020) **Application of Exogenous dsRNAs-induced RNAi in Agriculture: Challenges and Triumphs** *Front Plant Sci* **11** <https://doi.org/10.3389/fpls.2020.00946>
10. Devanapally S, Raman P, Chey M, Allgood S, Ettefa F, Diop M, Lin Y, Cho YE, Jose AM (2021) **Mating can initiate stable RNA silencing that overcomes epigenetic recovery** *Nat Commun* **12** <https://doi.org/10.1038/s41467-021-24053-4>
11. Dokshin GA, Ghanta KS, Piscopo KM, Mello CC (2018) **Robust Genome Editing with Short Single-Stranded and Long, Partially Single-Stranded DNA Donors in *Caenorhabditis elegans*** *Genetics* **210**:781–787 <https://doi.org/10.1534/genetics.118.301532>
12. Duchaine TF *et al.* (2006) **Functional Proteomics Reveals the Biochemical Niche of *C. elegans* DCR-1 in Multiple Small-RNA-Mediated Pathways** *Cell* **124**:343–354 <https://doi.org/10.1016/j.cell.2005.11.036>
13. Feinberg EH, Hunter CP (2003) **Transport of dsRNA into Cells by the Transmembrane Protein SID-1** *Science* **301**:1545–1547 <https://doi.org/10.1126/science.1087117>

14. Ferguson EL, Horvitz HR (1985) **IDENTIFICATION AND CHARACTERIZATION OF 22 GENES THAT AFFECT THE VULVAL CELL LINEAGES OF THE NEMATODE CAENORHABDITIS ELEGANS** *Genetics* **110**:17–72 <https://doi.org/10.1093/genetics/110.1.17>
15. Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC (1998) **Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*** *Nature* **391**:806–811 <https://doi.org/10.1038/35888>
16. Frøkjær-Jensen C, Davis MW, Ailion M, Jorgensen EM (2012) **Improved Mos1-mediated transgenesis in *C. elegans*** *Nat Methods* **9**:117–118 <https://doi.org/10.1038/nmeth.1865>
17. Gabel HW, Ruvkun G (2008) **The exonuclease ERI-1 has a conserved dual role in 5.8S rRNA processing and RNAi** *Nat Struct Mol Biol* **15**:531–533 <https://doi.org/10.1038/nsmb.1411>
18. Gu W *et al.* (2009) **Distinct Argonaute-Mediated 22G-RNA Pathways Direct Genome Surveillance in the *C. elegans* Germline** *Molecular Cell* **36**:231–244 <https://doi.org/10.1016/j.molcel.2009.09.020>
19. Guang S, Bochner AF, Pavelec DM, Burkhart KB, Harding S, Lachowiec J, Kennedy S (2008) **An Argonaute Transports siRNAs from the Cytoplasm to the Nucleus** *Science* **321**:537–541 <https://doi.org/10.1126/science.1157647>
20. Jobson MA, Jordan JM, Sandrof MA, Hibshman JD, Lennox AL, Baugh LR (2015) **Transgenerational Effects of Early Life Starvation on Growth, Reproduction, and Stress Resistance in *Caenorhabditis elegans*** *Genetics* **201**:201–212 <https://doi.org/10.1534/genetics.115.178699>
21. Jumper J *et al.* (2021) **Highly accurate protein structure prediction with AlphaFold** *Nature* **596**:583–589 <https://doi.org/10.1038/s41586-021-03819-2>
22. Kamath RS *et al.* (2003) **Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi** *Nature* **421**:231–237 <https://doi.org/10.1038/nature01278>
23. Kennedy S, Wang D, Ruvkun G (2004) **A conserved siRNA-degrading RNase negatively regulates RNA interference in *C. elegans*** *Nature* **427**:645–649 <https://doi.org/10.1038/nature02302>
24. Knight SW, Bass BL (2001) **A Role for the RNase III Enzyme DCR-1 in RNA Interference and Germ Line Development in *Caenorhabditis elegans*** *Science* **293**:2269–2271 <https://doi.org/10.1126/science.1062039>
25. Lee RC, Hammell CM, Ambros V (2006) **Interacting endogenous and exogenous RNAi pathways in *Caenorhabditis elegans*** *RNA* **12**:589–597 <https://doi.org/10.1261/rna.2231506>
26. Marré J, Traver EC, Jose AM (2016) **Extracellular RNA is transported from one generation to the next in *Caenorhabditis elegans*** *Proceedings of the National Academy of Sciences* **113**:12496–12501 <https://doi.org/10.1073/pnas.1608959113>
27. Matsumoto N, Sato K, Nishimasu H, Namba Y, Miyakubi K, Dohmae N, Ishitani R, Siomi H, Siomi MC, Nureki O (2015) **Crystal Structure and Activity of the Endoribonuclease Domain of the piRNA Pathway Factor Maelstrom** *Cell Reports* **11**:366–375 <https://doi.org/10.1016/j.celrep.2015.03.030>

28. Paix A, Folkmann A, Rasoloson D, Seydoux G (2015) **High Efficiency, Homology-Directed Genome Editing in *Caenorhabditis elegans* Using CRISPR-Cas9 Ribonucleoprotein Complexes** *Genetics* **201**:47–54 <https://doi.org/10.1534/genetics.115.179382>
29. Pak J, Fire A (2007) **Distinct Populations of Primary and Secondary Effectors During RNAi in *C. elegans*** *Science* **315**:241–244 <https://doi.org/10.1126/science.1132839>
30. Pak J, Maniar JM, Mello CC, Fire A (2012) **Protection from Feed-Forward Amplification in an Amplified RNAi Mechanism** *Cell* **151**:885–899 <https://doi.org/10.1016/j.cell.2012.10.022>
31. Parker GS, Eckert DM, Bass BL (2006) **RDE-4 preferentially binds long dsRNA and its dimerization is necessary for cleavage of dsRNA to siRNA** *RNA* **12**:807–818 <https://doi.org/10.1261/rna.2338706>
32. Parrish S, Fire A (2001) **Distinct roles for RDE-1 and RDE-4 during RNA interference in *Caenorhabditis elegans*** *RNA* **7**
33. Phillips CM, Montgomery TA, Breen PC, Ruvkun G (2012) **MUT-16 promotes formation of perinuclear Mutator foci required for RNA silencing in the *C. elegans* germline** *Genes Dev* **26**:1433–1444 <https://doi.org/10.1101/gad.193904.112>
34. Phillips CM, Updike DL (2022) **Germ granules and gene regulation in the *Caenorhabditis elegans* germline** *Genetics* **220** <https://doi.org/10.1093/genetics/iyab195>
35. Pinzón N, Bertrand S, Subirana L, Busseau I, Escrivá H, Seitz H (2019) **Functional lability of RNA-dependent RNA polymerases in animals** *PLOS Genetics* **15** <https://doi.org/10.1371/journal.pgen.1007915>
36. Preston MA, Porter DF, Chen F, Buter N, Lapointe CP, Keles S, Kimble J, Wickens M (2019) **Unbiased screen of RNA tailing activities reveals a poly(UG) polymerase** *Nat Methods* **16**:437–445 <https://doi.org/10.1038/s41592-019-0370-6>
37. Raman P, Zaghab SM, Traver EC, Jose AM (2017) **The double-stranded RNA binding protein RDE-4 can act cell autonomously during feeding RNAi in *C. elegans*** *Nucleic Acids Research* **45**:8463–8473 <https://doi.org/10.1093/nar/gkx484>
38. Ruby JG, Jan C, Player C, Axtell MJ, Lee W, Nusbaum C, Ge H, Bartel DP (2006) **Large-Scale Sequencing Reveals 21U-RNAs and Additional MicroRNAs and Endogenous siRNAs in *C. elegans*** *Cell* **127**:1193–1207 <https://doi.org/10.1016/j.cell.2006.10.040>
39. Sanan-Mishra N, Abdul Kader Jailani A, Mandal B, Mukherjee SK (2021) **Secondary siRNAs in Plants: Biosynthesis, Various Functions, and Applications in Virology** *Frontiers in Plant Science* **12**
40. Sandy P, Ventura A, Jacks T (2005) **Mammalian RNAi: a practical guide** *BioTechniques* **39**:215–224 <https://doi.org/10.2144/05392RV01>
41. Sapetschnig A, Sarkies P, Lehrbach NJ, Miska EA (2015) **Tertiary siRNAs Mediate Paramutation in *C. elegans*** *PLoS Genet* **11** <https://doi.org/10.1371/journal.pgen.1005078>
42. Seroussi U, Li C, Sundby AE, Lee TL, Claycomb JM, Saltzman AL (2022) **Mechanisms of epigenetic regulation by *C. elegans* nuclear RNA interference pathways** *Seminars in Cell & Developmental Biology* **127**:142–154 <https://doi.org/10.1016/j.semcd.2021.11.018>

43. Shukla A, Yan J, Pagano DJ, Dodson AE, Fei Y, Gorham J, Seidman JG, Wickens M, Kennedy S (2020) . **poly(UG)-tailed RNAs in genome protection and epigenetic inheritance** *Nature* **582**:283–288 <https://doi.org/10.1038/s41586-020-2323-8>
44. Sievers F *et al.* (2011) **Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega** *Molecular Systems Biology* **7** <https://doi.org/10.1038/msb.2011.75>
45. Sim G, Kehling AC, Park MS, Secor J, Divoky C, Zhang H, Malhotra N, Bhagdikar D, Abd El-Wahab EW, Nakanishi K (2022) **Manganese-dependent microRNA trimming by 3' → 5' exonucleases generates 14-nucleotide or shorter tiny RNAs** *Proceedings of the National Academy of Sciences* **119** <https://doi.org/10.1073/pnas.2214335119>
46. Steiner FA, Okihara KL, Hoogstrate SW, Sijen T, Ketting RF (2009) **RDE-1 slicer activity is required only for passenger-strand cleavage during RNAi in *Caenorhabditis elegans*** *Nat Struct Mol Biol* **16**:207–211 <https://doi.org/10.1038/nsmb.1541>
47. Stoyko D, Genzor P, Haase AD (2022) **Hierarchical length and sequence preferences establish a single major piRNA 3'-end** *iScience* **25** <https://doi.org/10.1016/j.isci.2022.104427>
48. Tabara H, Sarkissian M, Kelly WG, Fleenor J, Grishok A, Timmons L, Fire A, Mello CC (1999) **The rde-1 Gene** *RNA Interference, and Transposon Silencing in C. elegans*. *Cell* **99**:123–132 [https://doi.org/10.1016/S0092-8674\(00\)81644-X](https://doi.org/10.1016/S0092-8674(00)81644-X)
49. Tabara H, Yigit E, Siomi H, Mello CC (2002) **The dsRNA Binding Protein RDE-4 Interacts with RDE-1, DCR-1, and a DExH-Box Helicase to Direct RNAi in C. elegans** *Cell* **109**:861–871 [https://doi.org/10.1016/S0092-8674\(02\)00793-6](https://doi.org/10.1016/S0092-8674(02)00793-6)
50. Timmons L, Fire A (1998) **Specific interference by ingested dsRNA** *Nature* **395**:854–854 <https://doi.org/10.1038/27579>
51. Tsai H-Y, Chen C-CG, Conte D, Moresco JJ, Chaves DA, Mitani S, Yates JR, Tsai M-D, Mello CC (2015) **A Ribonuclease Coordinates siRNA Amplification and mRNA Cleavage during RNAi** *Cell* **160**:407–419 <https://doi.org/10.1016/j.cell.2015.01.010>
52. Uebel CJ, Anderson DC, Mandarino LM, Manage KI, Aynaszyan S, Phillips CM (2018) **Distinct regions of the intrinsically disordered protein MUT-16 mediate assembly of a small RNA amplification complex and promote phase separation of Mutator foci** *PLoS Genet* **14** <https://doi.org/10.1371/journal.pgen.1007542>
53. Varadi M *et al.* (2022) **AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models** *Nucleic Acids Research* **50**:D439–D444 <https://doi.org/10.1093/nar/gkab1061>
54. Vogel E, Santos D, Mingels L, Verdonck T-W, Broeck JV (2019) **RNA Interference in Insects: Protecting Beneficials and Controlling Pests** *Front Physiol* **9** <https://doi.org/10.3389/fphys.2018.01912>
55. Winston WM, Molodowitch C, Hunter CP (2002) **Systemic RNAi in C. elegans Requires the Putative Transmembrane Protein SID-1** *Science* **295**:2456–2459 <https://doi.org/10.1126/science.1068836>

56. Yang H, Zhang Y, Vallandingham J, Li H, Florens L, Mak HY (2012) **The RDE-10/RDE-11 complex triggers RNAi-induced mRNA degradation by association with target mRNA in *C. elegans*** *Genes Dev* **26**:846–856 <https://doi.org/10.1101/gad.180679.111>
57. Yigit E, Batista PJ, Bei Y, Pang KM, Chen C-CG, Tolia NH, Joshua-Tor L, Mitani S, Simard MJ, Mello CC (2006) **Analysis of the *C. elegans* Argonaute Family Reveals that Distinct Argonautes Act Sequentially during RNAi** *Cell* **127**:747–757 <https://doi.org/10.1016/j.cell.2006.09.033>
58. Zhang C, Montgomery TA, Fischer SEJ, Garcia SMDA, Riedel CG, Fahlgren N, Sullivan CM, Carrington JC, Ruvkun G (2012) **The *Caenorhabditis elegans* RDE-10/RDE-11 Complex Regulates RNAi by Promoting Secondary siRNA Amplification** *Current Biology* **22**:881–890 <https://doi.org/10.1016/j.cub.2012.04.011>
59. Zhang C, Montgomery TA, Gabel HW, Fischer SEJ, Phillips CM, Fahlgren N, Sullivan CM, Carrington JC, Ruvkun G (2011) . **mut-16 and other mutator class genes modulate 22G and 26G siRNA pathways in *Caenorhabditis elegans*** *Proc Natl Acad Sci USA* **108**:1201–1208 <https://doi.org/10.1073/pnas.1018695108>
60. Zhang D, Xiong H, Shan J, Xia X, Trudeau VL (2008) **Functional insight into Maelstrom in the germline piRNA pathway: a unique domain homologous to the DnaQ-H 3'–5' exonuclease, its lineage-specific expansion/loss and evolutionarily active site switch** *Biol Direct* **3** <https://doi.org/10.1186/1745-6150-3-48>
61. Zhang Y (2005) **TM-align: a protein structure alignment algorithm based on the TM-score** *Nucleic Acids Research* **33**:2302–2309 <https://doi.org/10.1093/nar/gki524>
62. Zhu Y, Zhu L, Wang X, Jin H (2022) **RNA-based therapeutics: an overview and prospectus** *Cell Death Dis* **13** <https://doi.org/10.1038/s41419-022-05075-2>
63. Zhuang JJ, Hunter CP (2011) **Tissue Specificity of *Caenorhabditis elegans* Enhanced RNA Interference Mutants** *Genetics* **188**:235–237 <https://doi.org/10.1534/genetics.111.127209>

Article and author information

Daphne R. Knudsen-Palmer

Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, USA.
Biological Sciences Graduate Program, University of Maryland, College Park, USA
ORCID iD: [0000-0001-5745-1185](https://orcid.org/0000-0001-5745-1185)

Pravrutha Raman

Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, USA.
Biological Sciences Graduate Program, University of Maryland, College Park, USA
ORCID iD: [0000-0002-7435-9425](https://orcid.org/0000-0002-7435-9425)

Farida Etefa

Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, USA.
Biological Sciences Graduate Program, University of Maryland, College Park, USA
ORCID iD: [0000-0003-0433-5951](https://orcid.org/0000-0003-0433-5951)

Laura De Ravin

Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, USA.
Biological Sciences Graduate Program, University of Maryland, College Park, USA

Antony M. Jose

Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, USA.
Biological Sciences Graduate Program, University of Maryland, College Park, USA

For correspondence: amjose@umd.edu

ORCID iD: [0000-0003-1405-0618](https://orcid.org/0000-0003-1405-0618)

Copyright

© 2024, Knudsen-Palmer et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Pedro Batista

National Institutes of Health, National Cancer Institute, Bethesda, United States of America

Senior Editor

Lori Sussel

University of Colorado Anschutz Medical Campus, Aurora, United States of America

Reviewer #1 (Public Review):

The goal of Knudsen-Palmer et al. was to define a biological set of rules that dictate the differential RNAi-mediated silencing of distinct target genes, motivated by facilitating the long-term development of effective RNAi-based drugs/therapeutics. To achieve this, the authors use a combination of computational modeling and RNAi function assays to reveal several criteria for effective RNAi-mediated silencing. This work provides insights into how (1) cis-regulatory elements influence the RNAi-mediated regulation of genes; (2) it is determined that genes can "recover" from RNAi-silencing signals in an animal; and (3) pUGylation occurs exclusively downstream of the dsRNA trigger sequence, suggesting 3° siRNAs are not produced. In addition, the authors show that the speed at which RNAi-silencing is triggered does not correlate with the longevity of the silencing. These insights are significant because they suggest that if we understand the rules by which RNAi pathways effectively silence genes with different transcription/processing levels then we can design more effective synthetic RNAi-based therapeutics targeting endogenous genes. The conclusions of this study are mostly supported by the data, but there are some aspects that need to be clarified.

(1) The methods do not describe the "aged RNAi plates feeding assay" in Figure 2E. The figure legend states that "aged RNAi plates" were used to trigger weaker RNAi, but the detail explaining the experiment is insufficient. How aged is aged? If the goal was to effectively reduce the dsRNA load available to the animals, why not quantitatively titrate the dsRNA provided? Were worms previously fed on the plates, or was simply a lawn of bacteria grown until presumably the IPTG on the plate was exhausted?

(2) Is the data presented in Figure 2F completed using the "aged RNAi plates" to achieve the partial silencing of *dpy-7* observed? Clarification of this point would be helpful.

(3) Throughout the manuscript the authors refer to "non-dividing cells" when discussing animals' ability to recover from RNA silencing. It is not clear what the authors specifically mean with the phrase "non-dividing cells", but as this is referred to in one of their major findings, it should be clarified. Do they mean the cells are somatic cells in aged animals, thus if they are "non-dividing" the siRNA pools within the cells cannot be diluted by cell division? Based on the methods, the animals of RNAi assays were L4/Young adults that were scored over 8 days after the initial pulse of dsRNA feeding. If this is the case, wouldn't these animals be growing into gravid adults after the feeding, and thus have dividing cells as they grew?

(4) What are the typical expression levels/turnover of *unc-22* and *bli-1*? Based on the results from the altered cis-regulatory regions of *bli-1* and *unc-22* in Figure 5, it seems like the transcription/turnover rates of each of these genes could also be used as a proof of principle for testing the model proposed in Figure 4. The strength of the model would be further increased if the RNAi sensitivity of *unc-22* reflects differences in its transcription/turnover rates compared to *bli-1*.

<https://doi.org/10.7554/eLife.97487.1.sa1>

Reviewer #2 (Public Review):

Summary:

This manuscript by Knudsen-Palmer et al. describes and models the contribution of MUT-16 and RDE-10 in the silencing through RNAi by the Argonaute protein NRDE-3 or others. The authors show that MUT-16 and RDE-10 constitute an intersecting network that can be redundant or not depending on the gene being targeted by RNAi. In addition, the authors provide evidence that increasing dsRNA processing can compensate for NRDE-3 mutants. Overall, the authors provide convincing evidence to understand the factors involved in RNAi in *C. elegans* by using a genetic approach.

Major Strengths:

The author's work presents a compelling case for understanding the intricacies of RNA interference (RNAi) within the model organism *Caenorhabditis elegans* through a meticulous genetic approach. By harnessing genetic manipulation, they delve into the role of MUT-16 and RDE-10 in RNAi, offering a nuanced understanding of the molecular mechanisms at play in two independent case study targets (*unc-22* and *bli-1*).

Major Weaknesses:

(1) It is unclear how the molecular mechanisms of amplification are different under the MUT-16 and RDE-10 branches of the regulatory pathway, since they are clearly distinct proteins structurally. It would be interesting to do some small-RNA-seq of products generated from *unc-22* and *bli-1*, on wild-type conditions and some of the mutants studied (eg. *mut-16*, *rde-10* and *mut-16 + rde-10*). That would provide some insights into whether the products of the 2 amplifications are the same in all conditions, just changing in abundance, or whether they are distinct in sequence patterns.

(2) In the same line, Figure 5 aims to provide insights into the sequence determinants that influence the RNAi of *bli-1*. It is unclear whether the changes in transcript stability dictated by the 3'UTR are the sole factor governing the preference for the MUT-16 and RDE-10 branches of the regulatory pathway. In line with the mutant *jam297*, it might be interesting to test whether factors like codon optimality, splicing, ... of the ORF region upstream from *bli-1*-dsRNA can affect its sensitivity to the MUT-16 and RDE-10 branches of the regulatory pathway.

SUPPLEMENTARY INFORMATION

Target-specific requirements for RNA interference can arise through restricted RNA amplification despite the lack of specialized pathways

Daphne R. Knudsen-Palmer¹, Pravrutha Raman^{1,2}, Farida Etefa^{1,3}, Laura De Ravin¹, and Antony M. Jose¹

¹Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, USA.
Biological Sciences Graduate Program, University of Maryland, College Park, USA.

²Current address: Division of Basic Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA, USA.

³Current address: Institute for Systems Genetics, New York University School of Medicine, New York, NY, USA.

*Corresponding author: Antony M. Jose

Email: amjose@umd.edu

This PDF file includes:

Figs. S1 to S7

Tables S1 to S3

SUPPLEMENTAL FIGURES AND TABLES





Fig. S1. A forward-genetic screen identifies a mutation that is expected to disrupt the Maelstrom domain of RDE-10. (A) Schematic depicting the screen. Males containing a transgene (*mex-5p::mCherry::h2b::tbb-2 3'utr::gpd-2 operon::gfp::h2b::cye-1 3'utr*) were mated to wild-type hermaphrodites, and progeny that showed silencing of the transgene were isolated. Animals that had been thus silenced for over 200 generations were mutagenized with N-ethyl-N-nitrosourea (ENU). Homozygous mutants that showed a re-activation of the transgene expression were isolated, and then subjected to a secondary feeding RNAi screen where the ability to silence two targets (*bli-1* and *unc-22*) was tested. (B) Schematics depicting the genes that were targeted by feeding RNAi. Black boxes, indicating exons, and regions that share homology with the dsRNA (blue, *bli-1*; orange, *unc-22*) are shown. (C) Predicted structure of RDE-10 showing the expected steric hindrance caused by the Ser228Phe mutation isolated in the screen. (*left*) The structure predicted by AlphaFold (Jumper et al., 2021; Varadi et al., 2022) contains a Maelstrom domain (teal; Pfam: PF13017). Ser228 (red) and the conserved EHC motif that can bind zinc (dark blue) are highlighted. (*middle* and *right*) Closeup of residue 228, showing the wild-type Ser (*middle*) and one orientation of Phe (*right*), which highlights the expected steric hindrance (red).

A

Alternative regulatory pathways:



Evidence against:


mut-16(-) mutants can silence *unc-22*

rde-10(-) mutants can silence *unc-22*

mut-16(-); nrde-3(-) mutants cannot silence *unc-22*

rde-10(-); nrde-3(-) mutants cannot silence *unc-22*

B



C




Fig. S2. Support for an intersecting regulatory network. (A) Alternative regulatory pathways excluded by data. Four plausible alternative models of the one presented in Fig. 2C along with the experimental evidence from Figs. 1 and 2 are shown. (B and C) Feeding RNAi of *unc-22* or *dpy-7*. Fractions silenced, numbers scored, comparisons, asterisks and error bars are as in Fig. 1. (B) An allele of *rde-10* that results in a premature stop codon (*jam243*) is still capable of silencing *unc-22* to near wild type levels. A newly created premature stop codon in *mut-16(-)* (*jam240*) created in the background of the RDE-10(Ser228Phe) mutant (*jam206*) fails to silence *unc-22*, supporting the findings in Fig. 1D. (C) An additional allele of *mut-16(-)* (*jam268*) shows about 25% silencing in response to *dpy-7* RNAi, supporting the findings in Fig. 2E.




Fig. S3. Overexpressing RDE-4 in the hypodermis. Semiquantitative RT-PCR of *rde-4* mRNA and *tbb-2* mRNA (control) in wild-type animals, *rde-4(-)* animals, or *rde-4(-)* animals expressing *rde-4(+)* in the hypodermis using a single-copy transgene (*Si(nas-9p)*). +RT and -RT indicate whether reverse transcriptase was used. The normalized mRNA abundance in *rde-4(-)* animals was subtracted from all lanes. With the observation in Fig. 3E and abundance of *rde-4* transcripts in the germline in wild-type animals (in situ data from NEXTDB), expression at ~30% of wild type is expected to be an overexpression in the hypodermis.




Fig. S4. An equilibrium model for RNAi. (A) Model highlighting different RNA species and their interactions during RNA interference. *Left*, Schematic showing downstream events upon processing of intracellular dsRNA. *Right*, Equations for the pre-mRNA ($[p]_i$) and mRNA ($[m]_i$) concentrations after RNAi in terms of the concentrations before RNAi ($[p]$ and $[m]$) and the length of the transcript (l_m). Rate constants for transcription ($k1$), binding constants for complex formation (1° siRNA-mRNA ($k2$), 2° siRNA-pre-mRNA ($k3$), and 2° siRNA-mRNA ($k4$), and rate constants for splicing followed by export out of the nucleus ($k5$) were considered. (B) Distributions of allowed values for all constants that are consistent with knockdown. One million random sets of values from 1 to 10 were assigned for all constants and the 790 sets that supported knockdown ($[m]_i < [m]$) and residual presence of pre-mRNA and mRNA ($[m]_i > 0$ and $[p]_i > 0$) were selected and plotted. Most values of $k2$ were small as expected to satisfy the $[m]_i > 0$ constraint because its square is subtracted in the equation for $[m]_i$. (C) RNAi can result in a variety of residual concentrations of pre-mRNA and mRNA. *Left*, Different ratios of residual mRNA and pre-mRNA

$([m]_i/[p]_i)$ are possible for a given length of the target mRNA (l_m : 0 to 1). *Middle* and *Right*, Impact of $[m]$ on $[m]_i$ and $[p]_i$. *Middle*, When the knockdown is poor (<10%), the concentrations of mRNA after RNAi ($[m]_i$) remain close to concentrations before RNAi ($[m]$) as expected for all levels of residual pre-mRNA ($[p]_i$) as evidenced by the graded levels of $[m]_i$ that mirror the graded levels of $[m]$. *Right*, When the knockdown is strong (>90%), the low concentrations of mRNA after RNAi ($[m]_i$) can be associated with a range of values for $[p]_i$. (D) RNAi can alter the ratios of pre-mRNA and mRNA. *Left*, Changes in the ratios of mRNA to pre-mRNA upon RNAi ($[m]/[p]$ before to $[m]_i/[p]_i$ after) are possible for a variety of extents of knockdown ($kd = [m]_i/[m]$: 0 to 1). *Middle* and *Right*, Impact of kd on $[m]/[p]$ versus $[m]_i/[p]_i$. Increases and decreases in the ratios of mRNA to pre-mRNA (above versus below the dotted line) can occur in response to both poor (<10% with $0.9 < kd < 1.0$, *middle*) and good (>90% with $0.0 < kd < 0.1$, *right*) knockdown. While it appears that most cases of efficient knockdown are associated with a decrease in the ratio of mRNA to pre-mRNA (i.e., only one point above dotted line in *right*), increasing the numbers of parameter sets explored could reveal additional examples. (E) (left) Same as in (D, *right*) except only values that were to the left and above the dotted line. Instead of one million simulated values, one hundred million values were simulated to better examine the range of possible values in this group. (*center*) Same as in (B) but for the values culled in (E, *left*). (F) Constants (k_1 to k_5) do not appear to show strong correlations with one another, except for k_2 and k_4 . The lack of values in the bottom left corner comparing k_2 to k_4 (bottom graphs) reflects the expectation that efficient silencing is not possible when both the primary and secondary binding constants are low.




Fig. S5. Impact of model parameters on the time to knockdown (τ_{kd}) and the duration of knockdown (t_{kd}). A low (blue) and high (red) value was selected for each parameter and the τ_{kd} and t_{kd} that result upon similarly varying all other parameters were plotted. Comparisons for 1^o siRNA production ((A), $k1 = 0.1$ vs. 1.0), recognition of transcripts and 1^o siRNAs ((B), $k2 = 0.01$ vs. 0.1), pUG RNA production ((C), $k3 = 0.1$ vs. 1.0), 2^o siRNA production ((D), $k4 = 0.1$ vs. 1.0), binding of transcripts and 2^o siRNAs ((E), $k5 = 0.01$ vs. 0.1), transcript maturation, i.e., splicing and export out of the nucleus ((F), $k6 = 0.01$ vs. 0.1), downregulation of transcription ((G), $k8 = 0.01$ vs. 0.1), length of dsRNA ((H), $l_{ds} = 220$ vs. 440), length of target mRNA ((I), $l_m = 1$ vs. 10), mRNA turnover ((J), $T_m = 0.005$ vs. 0.05), pre-mRNA turnover ((K), $T_p = 0.005$ vs. 0.05), 1^o siRNA turnover ((L), $T_{pri} = 0.005$ vs. 0.05), 2^o siRNA turnover ((M), $T_{sec} = 0.005$ vs. 0.05), and pUG RNA turnover ((N), $T_{ug} = 0.005$ vs. 0.05) are shown.




Fig.S6. Replacing the *unc-22* 3' cis-regulatory sequence with that of *bli-1* results in defects that show a NRDE-3-dependent enhancement when exposed to *unc-22*-dsRNA. (A) Schematic depicting the removal of the endogenous *unc-22* 3' cis-regulatory region and the addition of the *bli-1* 3' cis-regulatory region. (B) Silencing in response to *unc-22*-dsRNA (orange) or the control vector (grey). Fractions silenced, numbers scored, comparisons, asterisks and error bars are as in Fig. 1. Disruption of the *unc-22* 3' cis-regulatory region results in animals that show twitching in 3mM levamisole even without the addition of *unc-22* dsRNA. Addition of *unc-22*-dsRNA resulted in significantly more animals with altered cis-regulatory regions (*unc-22(jam300)*, *unc-22(jam301)*) showing twitching. This enhancement was not observed in animals that also lack NRDE-3 (*unc-22(jam300); nrde-3(jam205)*). Given the low effect size for NRDE-3 dependence, we do not interpret these results as conclusive, but rather present the data for completeness.

Table S1. Strains used in this study.

Strain name	Genotype
N2	wild type
AMJ174	<i>oxSi487[mex-5p::mCherry::H2B::tbb-2 3'UTR::gpd-2 operon::GFP::H2B::cye-1 3'UTR + unc-119(+)] dpy-2(jam29) II; unc-119(ed3)? III; lin-2(jam30) X</i>
AMJ183	<i>rde-4(ne301) III; nrde-3(tm1116) X</i>
AMJ285	<i>jamSi1 [mex-5p::rde-4(+)] II; rde-4(ne301) III</i>
AMJ345	<i>jamSi2 [mex-5p::rde-1(+)] II; rde-1(ne219) V</i>
AMJ422	<i>jamSi6 [nas-9p::rde-4(+)] II; unc-119(ed) III</i>
AMJ489	<i>nrde-3(tm1116) X; eri-1(mg366) IV</i>
AMJ565	<i>jamSi6 [Pnas-9::rde-4(+):rde-4 3'UTR] II; unc-119(ed3) III rde-4(ne301) III</i>
AMJ611	<i>jamSi6 [nas-9p::rde-4(+):rde-4 3'UTR] II; rde-4(ne301) III; nrde-3(tm1116) X</i>
AMJ1023	<i>mut-16(jam138) I; oxSi487 dpy-2(jam29) II; unc-119(ed3)? III; lin-2(jam30) X</i>
AMJ1025	<i>mut-16(jam139) rde-10(jam248) I; oxSi487 dpy-2(jam29) II; unc-119(ed3)? III; lin-2(jam30) X</i>
AMJ1035	<i>mut-16(jam140) I; oxSi487 dpy-2(jam29) II; unc-119(ed3)? III; lin-2(jam30) X</i>
AMJ1042	<i>mut-16(jam141) I; oxSi487 dpy-2(jam29) II; unc-119(ed3)? III; lin-2(jam30) X</i>
AMJ1091	<i>mut-16(jam247) I; oxSi487 dpy-2(jam29) II; unc-119(ed3)? III; lin-2(jam30) X</i>
AMJ1195	<i>jamSi59 [Pmex-5::gfp::cye-1 3'UTR + unc-119(+)] II; unc-119(ed3) III</i>
AMJ1397	<i>mut-16(jam148) I</i>
AMJ1470	<i>mut-16(jam148) rde-10(jam196) I</i>
AMJ1489	<i>rde-10(jam206) I</i>
AMJ1510	<i>nrde-3(jam205) X</i>
AMJ1545	<i>mut-16(jam148) I; nrde-3(jam205) X</i>
AMJ1568	<i>rde-10(jam206) I; nrde-3(jam205) X</i>
AMJ1611	<i>mut-16(jam240) rde-10(jam206) I</i>
AMJ1614	<i>rde-10(jam243) I</i>
AMJ1621	<i>eri-1(mg366) IV; nrde-3(jam205) X</i>
AMJ1622	<i>rde-10(jam206) I; eri-1(mg366) IV</i>
AMJ1623	<i>rde-10(jam206) I; eri-1(mg366) IV</i>
AMJ1624	<i>rde-10(jam206) I; eri-1(mg366) IV; nrde-3(jam205) X</i>
AMJ1625	<i>rde-10(jam206) I; eri-1(mg366) IV; nrde-3(jam205) X</i>
AMJ1631	<i>mut-16(jam148) I; eri-1(mg366) IV; nrde-3(jam205) X</i>
AMJ1632	<i>mut-16(jam148) I; eri-1(mg366) IV; nrde-3(jam205) X</i>
AMJ1657	<i>mut-16(jam240) rde-10(jam206) I; eri-1(jam260) IV</i>
AMJ1658	<i>mut-16(jam240) rde-10(jam206) I; eri-1(jam261) IV</i>
AMJ1659	<i>mut-16(jam240) rde-10(jam206) I; eri-1(jam262) IV</i>
AMJ1660	<i>eri-1(jam263) IV</i>
AMJ1661	<i>eri-1(jam264) IV</i>
AMJ1672	<i>mut-16(jam265) I; eri-1(jam263) IV</i>
AMJ1673	<i>mut-16(jam266) I; eri-1(jam264) IV</i>
AMJ1674	<i>mut-16(jam267) I; eri-1(jam264) IV</i>
AMJ1675	<i>mut-16(jam268) I</i>
AMJ1709	<i>jam297[bli-1p::gfp::cye-1 3'utr + unc-119(+)] II; unc-119(ed3) III</i>
AMJ1721	<i>mut-16(jam148) I; jamSi6 [nas-9p::rde-4(+)] II; unc-119(ed)? III</i>
AMJ1722	<i>mut-16(jam148) I; jamSi6 II; unc-119(ed)? III; eri-1(jam263) IV</i>
AMJ1723	<i>rde-10(jam206) I; jamSi6 II; unc-119(ed)? III</i>
AMJ1724	<i>rde-10(jam206) I; jamSi6 II; unc-119(ed)? III; eri-1(mg366) IV</i>
AMJ1725	<i>rde-10(jam206) I; jam297[bli-1p::gfp::cye-1 3'utr + unc-119(+)] II; unc-119(ed3)? III</i>
AMJ1726	<i>jam297[bli-1p::gfp::cye-1 3'utr + unc-119(+)] II; unc-119(ed3)? III; nrde-3(jam205) X</i>
AMJ1727	<i>mut-16(jam148) I; jam297[bli-1p::gfp::cye-1 3'utr + unc-119(+)] II; unc-119(ed3)? III</i>
AMJ1728	<i>mut-16(jam148) rde-10(jam196) I; jam297[bli-1p::gfp::cye-1 3'utr + unc-119(+)] II; unc-119(ed3)? III</i>

AMJ1730	<i>unc-22(jam300) IV</i>
AMJ1731	<i>unc-22(jam301) IV</i>
AMJ1754	<i>bli-1(jam307) II</i>
AMJ1755	<i>bli-1(jam308) II</i>
AMJ1757	<i>unc-22(jam300) IV; nrde-3(jam205) X</i>
AMJ1758	<i>bli-1(jam307) II; nrde-3(jam205) X</i>
AMJ1771	<i>bli-1(jam308) II; nrde-3(jam205) X</i>
EG6787	<i>oxSi487 II; unc-119(ed3) III</i>
GR1373	<i>eri-1(mg366) IV</i>
WM27	<i>rde-1(ne219) V</i>
WM49	<i>rde-4(ne301) III</i>
WM156	<i>nrde-3(tm1116) X</i>

Table S2. Oligonucleotides used in this study.

Primer	Sequence
P1	atttaggtgacactatagaaatgctcagagatgctcggttttagagctagaaatagcaag
P2	tcactttctcgtgcgttcc
P3	ggagaaccactcccagaatg
P4	aatcaatcggctgtccacac
P5	atttaggtgacactatagctggatcacctgggaatccggttttagagctagaaatagcaag
P6	aatcgcaaacgagtggttac
P7	cgggctagatcataatgagg
P8	ggaccacgtggagtccaggacatccagggtttccagggtgaccaggagagatggaatt
P9	gaatattttcgaaaata
P10	cggcacatgcgaatattttccgaaaatagaaggatatttcaactcgatccagaaaaac
P11	gctaccataggcaccgcatg
P12	cactgaaactcaatacggcaagatgagaatgactggaaaccgtaccgcatgcggtgcctatggtagcggagcttca catggctcagaccaacagccta
P13	cacaaacgccaggaaaggaag
P14	catttctgcgttggtggacc
P15	gttgaacggatatctctgc
P16	aagattgaatgttgaacgaatatttcagcaggatacgatgaaagcttattgattgatg
P17	ccgaaatccagatgagttcc
P18	gcatctggataaaaccaagc
P19	ccgatacaatcagaatgatc
P20	agcaaggccaccgatcaaatcagaatgattaggcagacaaggatattatgacaagatatt
P21	ggcattcgagccaataatgc
P22	cgttgctcggcaactct
P23	acaccacgtacaaatgtttg
P24	tgcgtcatccacaccacgtacaaacgttttagggcactgcaaaaaagccatccagccaaca
P25	gactgtgctgacgctgttt
P26	ctcccagtggtttcgttt
P27	tgtggacacggaatcagatc
P28	gaaacagtcgatgctgctccatatttccgataggatcttcaacggctgtacacatggatg
P29	cctatgtccgacctgtcaga
P30	caattccgatttctgaagag
P31	cagacctcacgatatgtggaaa
P32	ggaacatatggggcattcg
P33	caactttgatagaaaagt
P34	acaagttgtacaaaaagc
P35	gattacgccaagctatcaactttgatagaaaagttgcctaccaaagtagaaattcc
P36	acaactccagtagaaaagttcttctcttactcatgatgaggttagatcacacta
P37	tttcgctgcctgtcacactc
P38	tacgcggaagacccaaatg
P39	gaacgcgtcgaggtgatagc
P40	ataaggagttccacgcccag

P41 ctagtgagtcgtattataagtg
 P42 tgaagacgacgagccacttg
 P43 ctagaacttctcataatag
 P44 ggatacagagagaagcacaat
 P45 ctttacaggaaccactattatgagaagtttctagtttaacatctctgccaccaccact
 P46 ccttatctctgcggtttccaactctccgcttctccaacatttctcagtcaacag
 P47 catacagaaggagaaatcgc
 P48 gttgtagtacagtgtcgc
 P49 gcgtccaattctgaaatca
 P50 ggtggcaggatgattagaca
 P51 aattctcactcaaaatttgc
 P52 tgcaaaatagcggcagctcttctcctgtctaataactaaaaaaaacttctagtctaac
 P53 tgtctttcaaattctcactcaaaatttgcgtgatcgattggcttctctcgtatcc
 P54 gacgacgacggcatctatgt
 P55 gctatggctgttctcatggcggcgtcgccatattctacttcacacacacacacacaca
 P56 gctatggctgttctcatggc
 P57 gatttctacgatcacattct
 P58 tgctccgtggagcaactcgc
 P59 gagcacactattctgtgcat
 P60 ggcgtcgccatattctact
 P61 cacttgctggaaagacaagg
 P62 cgcaagcatgctggtttgta
 P63 gcattccatctgcaatgcga
 P64 cagtgtgctttaaactcggc
 P65 tgctcttcggcagttgcttc
 P66 gcaaagaatctgcagcatgg
 P67 gaacacaccagactgaaga
 P68 gacgagcaaagtctcaacg
 P69 tcgtcttcggcagttgcttc

Table S3. Summary of statistics.

Figure	Comparison	Total n	Silenced n	p value	Strains	notes
1B <i>bli-1</i>	wild type vs <i>mut-16(jam138)</i>	757, 229	391, 0	<0.00001, *	EG6787, AMJ1023	pooled EG6787 from separate experiments
1B <i>bli-1</i>	wild type vs <i>mut-16(jam140)</i>	757, 277	391, 0	<0.00001, *	EG6787, AMJ1035	pooled EG6787 from separate experiments
1B <i>bli-1</i>	wild type vs <i>mut-16(jam141)</i>	757, 124	391, 0	<0.00001, *	EG6787, AMJ1042	pooled EG6787 from separate experiments
1B <i>bli-1</i>	wild type vs <i>mut-16(jam247)</i>	757, 446	391, 1	<0.00001, *	EG6787, AMJ1091	pooled EG6787 from separate experiments
1B <i>bli-1</i>	wild type vs <i>mut-16(jam139) rde-10(jam248)</i>	757, 412	391, 1	<0.00001, *	EG6787, AMJ1025	pooled EG6787 from separate experiments
1B <i>unc-22</i>	wild type vs <i>mut-16(jam139) rde-10(jam248)</i>	309, 173	282, 0	<0.00001, *	EG6787, AMJ1025	pooled EG6787 from separate experiments
1B <i>unc-22</i>	<i>mut-16(jam138)</i> vs <i>mut-16(jam139) rde-10(jam248)</i>	180, 173	111, 0	<0.00001, *	AMJ1023, AMJ1025	pooled EG6787 from separate experiments
1B <i>unc-22</i>	<i>mut-16(jam140)</i> vs <i>mut-16(jam139) rde-10(jam248)</i>	91, 173	50, 0	<0.00001, *	AMJ1035, AMJ1025	pooled EG6787 from separate experiments

1B <i>unc-22</i>	mut-16(jam141) vs mut- 16(jam139) rde- 10(jam248)	267, 173	208, 0	<0.00001, *	AMJ1042, AMJ1025	pooled EG6787 from separate experiments
1B <i>unc-22</i>	mut-16(jam247) vs mut- 16(jam139) rde- 10(jam248)	100, 173	84, 0	<0.00001, *	AMJ1091, AMJ1025	pooled EG6787 from separate experiments
1D <i>bli-1</i>	wild type vs mut- 16(jam148)	202, 126	159, 0	<0.00001, *	N2, AMJ1397	
1D <i>bli-1</i>	wild type vs mut- 16(jam148) rde- 10(jam196)	202, 209	159, 0	<0.00001, *	N2, AMJ1470	
1D <i>bli-1</i>	wild type vs rde- 10(jam206)	202, 209	159, 0	<0.00001, *	N2, AMJ1489	
1D <i>unc-22</i>	wild type vs mut- 16(jam148)	206, 121	204, 119	0.58802, ns	N2, AMJ1397	
1D <i>unc-22</i>	wild type vs mut- 16(jam148) rde- 10(jam196)	206, 146	204, 0	<0.00001, *	N2, AMJ1470	
1D <i>unc-22</i>	wild type vs rde- 10(jam206)	206, 190	204, 185	0.21023, ns	N2, AMJ1489	
2B <i>bli-1</i>	wild type vs nrde-3(jam205)	295, 274	172, 16	<0.00001, *	N2, AMJ1510	
2B <i>bli-1</i>	wild type vs rde- 10(jam206); nrde-3(jam205)	295, 219	172, 0	<0.00001, *	N2, AMJ1568	
2B <i>bli-1</i>	wild type vs mut- 16(jam148); nrde-3(jam205)	295, 121	172, 0	<0.00001, *	N2, AMJ1545	
2B <i>unc-22</i>	wild type vs nrde-3(jam205)	129, 110	127, 108	0.87221, ns	N2, AMJ1510	
2B <i>unc-22</i>	wild type vs rde- 10(jam206); nrde-3(jam205)	129, 111	127, 0	<0.00001, *	N2, AMJ1568	
2B <i>unc-22</i>	wild type vs mut- 16(jam148); nrde-3(jam205)	129, 101	127, 0	<0.00001, *	N2, AMJ1545	
2E	wild type vs mut- 16(jam148)	183, 107	179, 5	<0.00001, *	N2, AMJ1397	
2E	wild type vs rde- 10(jam206)	183, 116	179, 16	<0.00001, *	N2, AMJ1489	
2E	wild type vs nrde-3(jam205)	183, 105	179, 72	<0.00001, *	N2, AMJ1510	
2E	mut-16(jam148) vs rde- 10(jam206)	107, 116	5, 16	0.0190241, *	AMJ1397, AMJ1489	
2E	rde-10(jam206) vs nrde- 3(jam205)	116, 105	16, 72	<0.00001, *	AMJ1489, AMJ1510	
2F <i>unc-54</i>	wild type vs mut- 16(jam148)	148, 124	142, 2	<0.00001, *	N2, AMJ1397	
2F <i>unc-54</i>	wild type vs rde- 10(jam206)	148, 128	142, 2	<0.00001, *	N2, AMJ1489	

2F <i>unc-54</i>	wild type vs nrde-3(jam205)	148, 263	142, 49	<0.00001, *	N2, AMJ1510	Fed on a different day with similar N2 silencing 164/171 animals showing Unc-54
2F <i>unc-54</i>	wild type vs mut-16(jam148) rde-10(jam196)	148, 159	142, 0	<0.00001, *	N2, AMJ1470	
2F <i>unc-54</i>	wild type vs mut-16(jam148); nrde-3(jam205)	148, 166	142, 0	<0.00001, *	N2, AMJ1545	Fed on a different day with similar N2 silencing 164/171 animals showing Unc-54
2F <i>unc-54</i>	wild type vs rde-10(jam206); nrde-3(jam205)	148, 206	142, 0	<0.00001, *	N2, AMJ1568	Fed on a different day with similar N2 silencing 164/171 animals showing Unc-54
2F <i>unc-54</i>	nrde-3(jam205) vs mut-16(jam148)	263, 124	49, 2	<0.00001, *	AMJ1510, AMJ1397	
2F <i>unc-54</i>	nrde-3(jam205) vs rde-10(jam206)	263, 128	49, 2	<0.00001, *	AMJ1510, AMJ1489	
2F <i>dpy-7</i>	wild type vs mut-16(jam148)	322, 130	279, 34	<0.00001, *	N2, AMJ1397	
2F <i>dpy-7</i>	wild type vs rde-10(jam206)	322, 159	279, 2	<0.00001, *	N2, AMJ1489	
2F <i>dpy-7</i>	wild type vs nrde-3(jam205)	322, 126	279, 3	<0.00001, *	N2, AMJ1510	
2F <i>dpy-7</i>	wild type vs mut-16(jam148) rde-10(jam196)	322, 184	279, 0	<0.00001, *	N2, AMJ1470	
2F <i>dpy-7</i>	wild type vs mut-16(jam148); nrde-3(jam205)	322, 99	279, 0	<0.00001, *	N2, AMJ1545	
2F <i>dpy-7</i>	wild type vs rde-10(jam206); nrde-3(jam205)	322, 180	279, 0	<0.00001, *	N2, AMJ1568	
2F <i>dpy-7</i>	mut-16(jam148) vs rde-10(jam206)	130, 159	34, 2	<0.00001, *	AMJ1397, AMJ1489	
2F <i>dpy-7</i>	mut-16(jam148) vs nrde-3(jam205)	130, 126	34, 6	<0.00001, *	AMJ1397, AMJ1510	
2F	wild type vs mut-16(jam148)	183, 107	179, 5	<0.00001, *	N2, AMJ1397	
2F	wild type vs rde-10(jam206)	183, 116	179, 16	<0.00001, *	N2, AMJ1489	
2F	wild type vs nrde-3(jam205)	183, 105	179, 72	<0.00001, *	N2, AMJ1510	
2F	mut-16(jam148) vs rde-10(jam206)	107, 116	5, 16	0.0190241, *	AMJ1397, AMJ1489	

2F	rde-10(jam206) vs nrde-3(jam205)	116, 105	16, 72	<0.00001, *	AMJ1489, AMJ1510	
3B	wild type vs eri-1(mg366)	200, 200	134, 145	0.231163, ns	N2, GR1373	N2 from a second experiment showed comparable values (219/304), GR1373 from a second experiment showed comparable values (169/213)
3B	nrde-3(jam205) vs eri-1(mg366); nrde-3(tm1116)	143, 200	3, 167	0.000132, *	N2, AMJ489	
3B	mut-16(148/268) vs mut-16(jam265-7); eri-1(jam263)	172, 354	0, 0	>0.5, ns	AMJ1397, AMJ1675, AMJ1672-4	AMJ1397 (0/110) and AMJ1675 (0/123) were pooled; AMJ1672 (0/130), AMJ1673 (0/88), and AMJ1674 (0/111) were pooled
3B	rde-10(jam206) vs rde-10(jam206); eri-1(mg366)	233, 329	0, 0	>0.5, ns	AMJ1489, AMJ1622, AMJ1623	AMJ1622 (0/171) and AMJ1623 (0/183) were pooled
3B	nrde-3(jam205) vs eri-1(mg366); nrde-3(jam205)	143, 169	3, 130	<0.00001, *	AMJ1510, AMJ1621	
3C	mut-16(jam148/240) rde-10(jam196/206) vs. mut-16(jam240) rde-10(jam 206); eri-1(jam260-2)	149, 191	1, 10	0.01826, *	AMJ1470, AMJ1611, AMJ1657-9	AMJ1470 (0/32) and AMJ1661 (1/116) were pooled; AMJ1657 (9/75), AMJ1658 (0/56), and AMJ1659 (1/50) were pooled
3C	rde-10(jam206); nrde-3(jam205) vs rde-10(jam206); eri-1(mg366); nrde-3(jam205)	113, 265	6, 18	0.58837, ns	AMJ1568, AMJ1624, AMJ1625	AMJ1624 (4/135) and AMJ1625 (14/112) were pooled
3C	mut-16(jam148); nrde-3(jam205) vs mut-16(jam148); eri-1(mg366); nrde-3(jam205)	28, 110	4, 1	0.00072, *	AMJ1545, AMJ1631, AMJ1632	AMJ1631 (1/52) and AMJ1632 (0/57) were pooled
3D <i>bli-1</i>	wild type vs rde-1(ne219)	50, 37	41, 0	<0.00001, *	N2, WM27	N2 data from <i>rde-4(-)</i> experiment; N2 data from <i>rde-1(-)</i> experiment (not shown) is comparable with 41/50 silenced

3D <i>bli-1</i>	wild type vs <i>rde-4(ne301)</i>	40, 50	40, 0	<0.00001, *	N2, WM49	N2 data from <i>rde-4(-)</i> experiment; N2 data from <i>rde-1(-)</i> experiment (not shown) is comparable with 41/50 silenced
3D <i>bli-1</i>	wild type vs <i>Si[mex-5p::rde-1(+)]/+; rde-1(ne219)</i>	50, 33	41, 0	<0.00001, *	N2, AMJ345	N2 data from <i>rde-4(-)</i> experiment; N2 data from <i>rde-1(-)</i> experiment (not shown) is comparable with 41/50 silenced
3D <i>bli-1</i>	wild type vs <i>Si[mex-5p::rde-4(+)]/+; rde-4(ne301)</i>	40, 41	40, 41	>0.5, ns	N2, AMJ285	N2 data from <i>rde-4(-)</i> experiment; N2 data from <i>rde-1(-)</i> experiment (not shown) is comparable with 41/50 silenced
3D <i>unc-22</i>	wild type vs <i>rde-1(ne219)</i>	25, 25	25, 0	<0.00001, *	N2, WM27	
3D <i>unc-22</i>	wild type vs <i>rde-4(ne301)</i>	25, 25	25, 0	<0.00001, *	N2, WM49	
3D <i>unc-22</i>	wild type vs <i>Si[mex-5p::rde-1(+)]/+; rde-1(ne219)</i>	25, 24	25, 2	<0.00001, *	N2, AMJ345	
3D <i>unc-22</i>	wild type vs <i>Si[mex-5p::rde-4(+)]/+; rde-4(ne301)</i>	25, 24	25, 16	0.001600, †	N2, AMJ285	† indicates statistical significance vs. wild type and vs. <i>rde-4(ne301)</i> mutant
3D <i>unc-22</i>	<i>rde-4(ne301)</i> vs <i>Si[mex-5p::rde-4(+)]/+; rde-4(ne301)</i>	25, 24	0, 16	<0.00001, †	WM49, AMJ285	† indicates statistical significance vs. wild type and vs. <i>rde-4(ne301)</i> mutant
3E	wild type vs <i>rde-4(ne301); nrde-3(tm1116)</i>	150, 200	142, 0	<0.00001, *	N2, AMJ183	
3E	wild type vs <i>rde-4(ne301); nrde-3(tm1116); Si[nas-9p::rde-4(+)]</i>	150, 150	142, 98	<0.00001, †	N2, AMJ611	† indicates statistical significance vs. wild type and vs. <i>rde-4(ne301); nrde-3(tm1116)</i> mutant
3E	<i>rde-4(ne301); nrde-3(tm1116); vs rde-4(ne301); nrde-3(tm1116); Si[nas-9p::rde-4(+)]</i>	200, 150	0, 98	<0.00001, †	AMJ183, AMJ611	† indicates statistical significance vs. wild type and vs. <i>rde-4(ne301); nrde-3(tm1116)</i> mutant
3F	wild type vs <i>eri-1(mg366); nrde-3(jam205)</i>	282, 140	85, 69	0.0001199, *	N2, AMJ1621	

3F	wild type vs rde-10(jam206); eri-1(mg366)	282, 279	85, 0	<0.00001, *	N2, AMJ1622	
3F	wild type v Si[nas-9p::rde-4(+)]; rde-10(jam206)	282, 180	85, 6	<0.00001, *	N2, AMJ1723	
3F	wild type v Si[nas-9p::rde-4(+)]; rde-10(jam206); eri-1(mg366)	282, 202	85, 0	<0.00001, *	N2, AMJ1724	
3F	wild type vs mut-16(jam148); eri-1(mg366)	282, 70	85, 0	<0.00001, *	N2, AMJ1672	
3F	wild type v Si[nas-9p::rde-4(+)]; mut-16(jam148)	282, 152	85, 0	<0.00001, *	N2, AMJ1721	
3F	wild type v Si[nas-9p::rde-4(+)]; mut-16(jam148); eri-1(mg366)	282, 187	85, 0	<0.00001, *	N2, AMJ1722	
5B	wild type vs bli-1(jam307)	488, 443	390, 425	<0.00001, *	N2, AMJ1754	
5B	wild type vs bli-1(jam308)	488, 245	390, 185	0.170996, ns	N2, AMJ1755	
5B	wild type vs nrde-3(jam205)	488, 377	390, 111	<0.00001, *	N2, AMJ1510	
5B	bli-1(jam307) vs bli-1(jam307); nrde-3(jam205)	443, 459	425, 313	<0.00001, *	AMJ1754, AMJ1758	
5B	nrde-3(jam205) vs bli-1(jam307); nrde-3(jam205)	377, 459	111, 313	<0.00001, *	AMJ1510, AMJ1758	
5B	bli-1(jam308) vs bli-1(jam308); nrde-3(jam205)	245, 117	185, 38	<0.00001, *	AMJ1755, AMJ1771	
5B	wild type vs bli-1(jam307)	488, 443	390, 425	<0.00001, *	N2, AMJ1754	
5D	<i>bli-1p::gfp</i> vs <i>mut-16(jam148)</i> ; <i>bli-1p::gfp</i>	21, 25	21, 6	<0.00001, *	AMJ1709, AMJ1727	
5D	<i>bli-1p::gfp</i> vs <i>rde-10(jam206)</i> ; <i>bli-1p::gfp</i>	21, 25	21, 9	<0.00001, *	AMJ1709, AMJ1725	
5D	<i>bli-1p::gfp</i> vs <i>nrde-3(jam205)</i> ; <i>bli-1p::gfp</i>	21, 25	21, 23	0.18508, ns	AMJ1709, AMJ1726	
5D	<i>bli-1p::gfp</i> vs <i>mut-16(jam148)</i> ; <i>rde-10(jam196)</i> ; <i>bli-1p::gfp</i>	21, 20	21, 0	<0.00001, *	AMJ1709, AMJ1728	

6A	1 vs 2 days post RNAi	94, 91	76, 86	0.00489, *	N2	
6A	1 vs 3 days post RNAi	94, 92	76, 89	0.00062, *	N2	
6A	1 vs 4 days post RNAi	94, 93	76, 89	0.001627, *	N2	
6A	5 days post RNAi versus 7 days on RNAi	110, 55	63, 51	<0.00001, *	N2	
6A	6 days post RNAi versus 7 days on RNAi	104, 55	61, 51	<0.00001, *	N2	
6A	7 days post RNAi versus 7 days on RNAi	112, 55	60, 51	<0.00001, *	N2	
6A	8 days post RNAi versus 7 days on RNAi	124, 55	45, 51	<0.00001, *	N2	
S2B	wild type vs rde-10(jam243)	115, 102	115, 100	0.13140071, ns	N2, AMJ1614	
S2B	wild type vs mut-16(jam240) rde-10(jam206)	115, 137	115, 5	<0.00001, *	N2, AMJ1611	
S2C	wild type vs mut-16(jam268)	322, 74	279, 20	<0.00001, *	N2, AMJ1675	From the same feed in Fig. 2E
S6B	wild type unc-22 vs wild type pL4440	138, 116	138, 0	<0.00001, *	N2	
S6B	wild type unc-22 vs nrde-3(jam205) unc-22	138, 110	138, 79	<0.00001, *	N2, AMJ1510	
S6B	unc-22(jam300) unc-22 vs unc-22(jam300) pL4440	101, 113	101, 101	0.00128, *	AMJ1730	
S6B	unc-22(jam301) unc-22 vs unc-22(jam301) pL4440	101, 65	101, 49	<0.00001, *	AMJ1731	
S6B	unc-22(jam300); nrde-3(jam205) unc-22 vs unc-22(jam300); nrde-3(jam205) pL4440	82, 108	70, 83	0.14214, ns	AMJ1757	
S6B	unc-22(jam300); nrde-3(jam205) unc-22 vs unc-22(jam300) unc-22	82, 101	70, 101	<0.00001, *	AMJ1757, AMJ1730	
S6B	unc-22(jam300); nrde-3(jam205) unc-22 vs nrde-	82, 110	70, 79	0.02592, *	AMJ1757, AMJ1510	

	3(jam205) unc-22					
--	------------------	--	--	--	--	--