

Instructor: Dr. Zhongchi Liu

3236 H.J. Patterson Hall

Phone: 5-1586

ZhongchiL@gmail.com

www.life.umd.edu/CBMG/faculty/liu/liu.html

TA: Jennifer Baxter

2114 (Lab) or 2114 (Office) Microbiology Building

Phone: 5-1758

Sugarpucker@gmail.com

About the course: www.life.umd.edu/classroom/BSCI410-Liu/BSCI410/

Lecture 1: Genes and Mutations

- Central dogma
- Genetic code
- Mutation type

Read: Ch 7 p191-193;
Ch 8 p238-240; 245-246

Figs: 7.2; 8.2; 8.3; 8.9

Reviewing basic terminology

Gene: a specific sequence of nucleotide pairs in a discrete region of DNA that acts as a functional unit, usually by encoding the instructions for making a particular protein

Transcription (where)

Translation (where)

Central dogma

Codon

ribosome

tRNA, rRNA, mRNA

Sense and antisense

- 5' to 3' direction of mRNA corresponds to N-terminal-to-C-terminal direction of polypeptide
 - One strand of DNA is a template
 - The other is an RNA-like strand
- Nonsense codons cause termination of a polypeptide chain
 - UAA (ocher), UAG (amber), and UGA (opal)




Fig. 8.9

**The Genetic Code: 61 triplet codons represent 20 amino acids;
3 triplet codons signify stop**

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

		Second letter				
		U	C	A	G	
First letter	U	UUU } Phe UUC } UUA } Leu UUG }	UCU } UCC } Ser UCA } UCG }	UAU } Tyr UAC } UAA Stop UAG Stop	UGU } Cys UGC } UGA Stop UGG Trp	U C A G
	C	CUU } CUC } Leu CUA } CUG }	CCU } CCC } Pro CCA } CCG }	CAU } His CAC } CAA } Gln CAG }	CGU } CGC } Arg CGA } CGG }	U C A G
	A	AUU } AUC } Ile AUA } AUG Met	ACU } ACC } Thr ACA } ACG }	AAU } Asn AAC } AAA } Lys AAG }	AGU } Ser AGC } AGA } Arg AGG }	U C A G
	G	GUU } GUC } Val GUA } GUG }	GCU } GCC } Ala GCA } GCG }	GAU } Asp GAC } GAA } Glu GAG }	GGU } GGC } Gly GGA } GGG }	U C A G

Summary of genetic code

- Codon consist of a triplet codon each of which specifies an amino acid
 - Code shows a 5' to 3' direction
- Codons are nonoverlapping
- Code includes three stop codons, UAA, UAG, and UGA that terminate translation
- Code is degenerate
- Fixed starting point establishes a reading frame
 - AUG in an initiation codon which specifies reading frame
- 5'- 3' direction of mRNA corresponds with N-terminus to C-terminus of polypeptide

Genotype vs. Phenotype

Phenotype depends on structure and abundance of a protein

Mutations: change in DNA, heritable, mostly devastating, few good. They alter protein structure and abundance, leading to changes of phenotype.

Alleles: different variations (genotypes) in a gene of different individuals