

distribution of cells free in flow. We analysed enough videotape to obtain 100 to 1,600 tethering events, and plotted the natural log of the number of cells that remained bound as a function of time after initiation of tethering. The most rapidly dissociating 90% or more of tethered cells were used to determine k_{off} .

Received 23 June 1997; accepted 13 February 1998.

1. Bell, G. I. Models for the specific adhesion of cells to cells: a theoretical framework for adhesion mediated by reversible bonds between cell surface molecules. *Science* **200**, 618–627 (1978).
2. Dembo, M., Torney, D. C., Saxman, K. & Hammer, D. The reaction-limited kinetics of membrane-to-surface adhesion and detachment. *Proc. R. Soc. Lond. B* **234**, 55–83 (1988).
3. Leckband, D., Müller, W., Schmitt, F.-J. & Ringsdorf, H. Molecular mechanisms determining the strength of receptor-mediated intermembrane adhesion. *Biophys. J.* **69**, 1162–1169 (1995).
4. Bruinsma, R. Les liaisons dangereuses: adhesion molecules do it statistically. *Proc. Natl Acad. Sci. USA* **94**, 375–376 (1997).
5. Evans, E. & Ritchie, K. Dynamic strength of molecular adhesion bonds. *Biophys. J.* **72**, 1541–1555 (1997).
6. Finger, E. B. *et al.* Adhesion through L-selectin requires a threshold hydrodynamic shear. *Nature* **379**, 266–269 (1996).
7. Alon, R., Fuhlbrigge, R. C., Finger, E. B. & Springer, T. A. Interactions through L-selectin between leukocytes and adherent leukocytes nucleate rolling adhesions on selectins and VCAM-1 in shear flow. *J. Cell Biol.* **135**, 849–865 (1996).
8. Lawrence, M. B., Kansas, G. S., Kunkel, E. J. & Ley, K. Threshold levels of fluid shear promote leukocyte adhesion through selectins (CD62L, P.E.). *J. Cell Biol.* **136**, 717–727 (1997).
9. Taylor, A. D., Neelamegham, S., Hellums, J. D., Smith, C. W. & Simon, S. I. Molecular dynamics of the transition from L-selectin to β_2 -integrin-dependent neutrophil adhesion under defined hydrodynamic shear. *Biophys. J.* **71**, 3488–3500 (1996).
10. Lasky, L. A. & Rosen, S. D. In *Inflammation: Basic Principles and Clinical Correlates* Vol. 2 (eds Gallin, J. I., Goldstein, I. J. & Snyderman, R.) 407–439 (Raven, New York, 1992).
11. Kansas, G. S. Selectins and their ligands: current concepts and controversies. *Blood* **88**, 3259–3287 (1996).
12. Springer, T. A. Traffic signals on endothelium for lymphocyte recirculation and leukocyte emigration. *Annu. Rev. Physiol.* **57**, 827–872 (1995).
13. Alon, R., Chen, S., Puri, K. D., Finger, E. B. & Springer, T. A. The kinetics of L-selectin tethers and the mechanics of selectin-mediated rolling. *J. Cell Biol.* **138**, 1169–1180 (1997).
14. Alon, R., Hammer, D. A. & Springer, T. A. Lifetime of the P-selectin: carbohydrate bond and its response to tensile force in hydrodynamic flow. *Nature* **374**, 539 (1995).
15. Norgard, K. E. *et al.* Enhanced interaction of L-selectin with the high endothelial venule ligand via selectively oxidized sialic acids. *Proc. Natl Acad. Sci. USA* **90**, 1068–1072 (1993).
16. Puri, K. D. & Springer, T. A. A Schiff base with mildly oxidized carbohydrate ligands stabilizes L-selectin and not P-selectin or E-selectin rolling adhesions in shear flow. *J. Biol. Chem.* **271**, 5404–5413 (1996).
17. Baumhueter, S. *et al.* Binding of L-selectin to the vascular sialomucin, CD34. *Science* **262**, 436–438 (1993).
18. Puri, K. D., Finger, E. B., Gaudernack, G. & Springer, T. A. Sialomucin CD34 is the major L-selectin ligand in human tonsil high endothelial venules. *J. Cell Biol.* **131**, 261–270 (1995).
19. Nicholson, M. W., Barclay, A. N., Singer, M. S., Rosen, S. D. & van der Merwe, P. A. Affinity and kinetic analysis of L-selectin (CD62L) binding to GlyCAM-1. *J. Biol. Chem.* **273**, 763–770 (1998).
20. Pierres, A., Benoliel, A.-M. & Bongrand, P. Experimental study of the rate of bond formation between individual receptor-coated spheres and ligand-bearing surfaces. *J. Phys. III France* **6**, 807–824 (1996).
21. Kyte, J. *Structure in Protein Chemistry* (Garland, New York, 1995).
22. Dembo, M. In *Some Mathematical Problems in Biology: Lectures on Mathematics in the Life Sciences* Vol. 25 (eds Goldstein, B. & Wofsy, C.) 1–27 (American Mathematical Society, Providence, 1994).
23. Puri, K. D., Finger, E. B. & Springer, T. A. The faster kinetics of L-selectin than of E-selectin and P-selectin rolling at comparable binding strength. *J. Immunol.* **158**, 405–413 (1997).
24. Chen, S., Alon, R., Fuhlbrigge, R. C. & Springer, T. A. Rolling and transient tethering of leukocytes on antibodies reveal specializations of selectins. *Proc. Natl Acad. Sci. USA* **94**, 3172–3177 (1997).

Acknowledgements. We thank T. K. Kishimoto, E. Butcher and G. Gaudernack for reagents.

Correspondence and requests for materials should be addressed to T.A.S.

Calcium oscillations increase the efficiency and specificity of gene expression

Ricardo E. Dolmetsch*†, Keli Xu* & Richard S. Lewis

Department of Molecular and Cellular Physiology, Beckman Center for Molecular and Genetic Medicine, Stanford University School of Medicine, Stanford, California 94305, USA

* These authors contributed equally to the work.

Cytosolic calcium ($[Ca^{2+}]_i$) oscillations are a nearly universal mode of signalling in excitable and non-excitable cells^{1–4}. Although Ca^{2+} is known to mediate a diverse array of cell functions, it is not known whether oscillations contribute to the efficiency or specificity of signalling or are merely an inevitable

consequence of the feedback control of $[Ca^{2+}]_i$. We have developed a Ca^{2+} clamp technique to investigate the roles of oscillation amplitude and frequency in regulating gene expression driven by the proinflammatory transcription factors NF-AT, Oct/OAP and NF- κ B. Here we report that oscillations reduce the effective Ca^{2+} threshold for activating transcription factors, thereby increasing signal detection at low levels of stimulation. In addition, specificity is encoded by the oscillation frequency: rapid oscillations stimulate all three transcription factors, whereas infrequent oscillations activate only NF- κ B. The genes encoding the cytokines interleukin (IL)-2 and IL-8 are also frequency-sensitive in a way that reflects their degree of dependence on NF-AT versus NF- κ B. Our results provide direct evidence that $[Ca^{2+}]_i$ oscillations increase both the efficacy and the information content of Ca^{2+} signals that lead to gene expression and cell differentiation.

Oscillations in $[Ca^{2+}]_i$ may be advantageous for receptor-mediated signal transduction, for example by increasing the fidelity of low-level signalling, preventing desensitization, or increasing signalling specificity^{1–4}; however, it has been difficult to demonstrate these and other possible functions⁵ for two reasons. First, the amplitude and frequency of $[Ca^{2+}]_i$ oscillations triggered through surface receptors varies among cells and in single cells over time, resulting in a mixture of stimulus waveforms that complicates analysis. Second, surface receptors are often coupled to multiple signalling pathways, making it difficult to ascribe downstream effects to Ca^{2+} alone. We have therefore developed a 'calcium clamp' method for generating homogeneous and synchronous receptor-independent $[Ca^{2+}]_i$ oscillations in large populations of T lymphocytes⁶ (Fig. 1). Ca^{2+} signals leading to T-cell activation are normally generated by a cascade involving antigen binding to the T-cell antigen receptor (TCR), generation of the second messenger inositol 1,4,5-trisphosphate ($InsP_3$), release of Ca^{2+} from internal stores, and Ca^{2+} influx across the plasma membrane⁷. Here we bypass the TCR/ $InsP_3$ pathway by treating Jurkat T cells with thapsigargin, an inhibitor of endoplasmic reticulum Ca^{2+} -ATPases that depletes internal Ca^{2+} stores and irreversibly activates store-operated Ca^{2+} (CRAC) channels in the plasma membrane⁸. Application of Ca^{2+} to cells treated in this way elevates $[Ca^{2+}]_i$ owing to influx through CRAC channels, whereas removal of extracellular Ca^{2+} allows pumps in the plasma membrane to return $[Ca^{2+}]_i$ to baseline levels. Thus, by rapidly changing the concentration of extracellular Ca^{2+} , it is possible to generate $[Ca^{2+}]_i$ oscillations having a uniform frequency across cells and an amplitude that is relatively constant in each cell over time (s.d., 12.7%) and among cells in the population (s.d., 18.2%; $n = 256$). A further advantage is that this technique probably mimics naturally occurring subcellular gradients of $[Ca^{2+}]_i$, because $[Ca^{2+}]_i$ oscillations triggered through the TCR result from periodic activation of CRAC channels^{9,10}.

We investigated whether oscillations affect the efficiency with which Ca^{2+} signals are detected. NF-AT is a Ca^{2+} -dependent transcription factor expressed in many cells, including T lymphocytes, in which it helps to regulate several immune-response genes including IL-2, IL-4 and tumour-necrosis factor- α (TNF- α)^{11,12}. NF-AT is activated by Ca^{2+} -stimulated dephosphorylation and translocation of a cytoplasmic subunit, which binds to a nuclear subunit induced by protein kinase C or by stimulation of the MAP kinase pathway^{11,12}. We compared the activity of an NF-AT/*lacZ* reporter gene¹³ in Jurkat cells stimulated with an oscillatory or constant elevation of $[Ca^{2+}]_i$ for 3 hours in the presence of 50 nM phorbol-12,13-dibutyrate (PdBu). Oscillations were generated with a period of 100 s, which is similar to the period in intact cells stimulated through the TCR^{9,14}. Oscillation amplitude was adjusted to produce the same average $[Ca^{2+}]_i$ as in the constant- $[Ca^{2+}]_i$ control cells to determine whether the kinetic features of the oscillations confer any signalling advantage relative to a sustained $[Ca^{2+}]_i$ increase, independently of the amount of Ca^{2+} that enters a cell. Figure 2a shows constant and oscillatory $[Ca^{2+}]_i$ stimuli in one

† Present address: Division of Neuroscience, Children's Hospital, Harvard University, 300 Longwood Avenue, Boston, Massachusetts 02120, USA.

experiment in which the average $[Ca^{2+}]_i$ was ~ 225 nM; subsequent fluorescence-activated cell sorting (FACS) analysis indicated that 39% of the cells stimulated with $[Ca^{2+}]_i$ oscillations were *lacZ*⁺, whereas only 19% of the cells with constant $[Ca^{2+}]_i$ were *lacZ*⁺. This enhancement of NF-AT-dependent transcription was not evident at higher $[Ca^{2+}]_i$ levels. In cells stimulated to an average of ~ 350 nM Ca^{2+} , oscillations activated roughly the same amount of *lacZ* expression as constant $[Ca^{2+}]_i$ elevation (Fig. 2b). The results of 12 paired experiments show that the induction of *lacZ* by constant $[Ca^{2+}]_i$ elevation falls off sharply below ~ 300 nM $[Ca^{2+}]_i$, whereas induction by oscillations is approximately constant down to at least 200 nM (Fig. 2c). The enhancement of NF-AT-dependent gene expression by oscillations thus increases as average $[Ca^{2+}]_i$ decreases below ~ 300 nM (Fig. 2d). In this way, oscillations appear to increase the ability of small Ca^{2+} signals to activate NFAT.

A long-standing question in cell signalling is how Ca^{2+} , with its abundant and varied intracellular targets, is able to achieve specificity and activate only a subset of those targets. One explanation is that variations in oscillation amplitude or frequency may discriminate among different Ca^{2+} -activated signalling pathways. In T cells, several transcription factors in addition to NF-AT are Ca^{2+} -sensitive, including NF- κ B and Oct/OAP^{15,16}. Both factors are involved in the control of IL-2 transcription, and NF- κ B also helps to regulate a large number of genes encoding cytokines, growth factors, adhesion molecules and other surface proteins^{11,17}. To determine whether the amplitude of the Ca^{2+} signal can distinguish among transcriptional pathways, we compared the Ca^{2+} sensitivity of luciferase reporter genes driven by NF-AT, Oct/OAP or NF- κ B in Jurkat cells. Phorbol ester (50 nM PdBu) or increased $[Ca^{2+}]_i$ (>600 nM) alone did not activate NF-AT or Oct/OAP measurably in thapsigargin-treated cells, and only slightly stimulated NF- κ B-dependent expression (~ 2.5 -fold above background). In contrast, the two stimuli acted together in a synergistic fashion to activate NF-AT, Oct/OAP and NF- κ B by 150-, 31- and 16-fold, respectively. In each case, the Ca^{2+} -dependent component of the response was completely suppressed by cyclosporin A, a potent and selective inhibitor of calcineurin,

consistent with a role for this phosphatase in mediating the effects of Ca^{2+} on these factors^{15,18,19}. As shown in Fig. 3a, all three pathways show a similar and highly nonlinear dependence on steady-state $[Ca^{2+}]_i$. The pronounced nonlinearity of NF-AT activation is consistent with previous studies^{5,20} and is an important determinant of the effects of oscillations on NF-AT signaling efficiency (see below). However, the similar Ca^{2+} dependencies of the three transcription factors suggests that oscillation amplitude is not likely to contribute significantly to selectivity among these pathways in T cells. In a previous study of B cells²¹, nuclear translocation of NF-AT appeared to be more Ca^{2+} -sensitive than that of NF- κ B; however, the transcriptional activity of these factors was not tested.

As oscillation amplitude is unlikely to discriminate among these three pathways, we tested whether oscillation frequency could. The calcium clamp was used to generate high-amplitude oscillations (~ 1 μ M peak) with periods from 100 to 1,800 s. As shown in Fig. 3b, the activation of all three transcription factors declines with increasing period. However, NF- κ B activation extends to oscillation periods as long as 1,800 s, in contrast to NF-AT and Oct/OAP, whose activity vanishes at periods ≥ 400 s. Frequency dependence of NF-AT has also been observed in response to photolytically generated pulses of $InsP_3$ (ref. 22), indicating that this behaviour is not specific to the use of thapsigargin to deplete stores. The difference in frequency requirements among the three factors is consistent with previous findings that NF- κ B persists in the nucleus for >16 min following a brief Ca^{2+} spike²¹, whereas NF-AT quickly exits following its rapid rephosphorylation^{21,23}. Thus, oscillation frequency dictates which combination of transcription factors is active; low frequencies recruit NF- κ B alone, whereas high frequencies activate NF-AT, Oct/OAP and NF- κ B. These results indicate that $[Ca^{2+}]_i$ oscillation frequency can discriminate among different transcriptional pathways.

The distinct frequency sensitivities of NF-AT, Oct/OAP and NF- κ B suggest that endogenous genes under the control of these factors may also respond differentially to oscillation frequency. We therefore compared the frequency sensitivities of genes encoding IL-2, an




Figure 1 Generation of synchronized $[Ca^{2+}]_i$ oscillations in T cell populations. **a**, Diagram of the Ca^{2+} clamp (see Methods). Oscillations are generated by alternately exposing cells with depleted stores to 0 or 1.5 mM Ca^{2+} . **b**, **c**, Average $[Ca^{2+}]_i$ in 200–300 cells in which the amplitude or frequency of oscillations was

varied. In **b**, Ca^{2+} was applied for 10 s (top), 15 s (middle), or 30 s (bottom) every 100 s. In **c**, Ca^{2+} was applied for 30-s periods with a interspike interval of 900 s (top), 400 s (middle), or 100 s (bottom).

essential growth factor for T cells, and IL-8, a chemokine involved in recruiting immune cells to sites of infection. Both genes can be activated in T cells by Ca^{2+} ionophore and phorbol ester^{24,25}, but their promoters are driven by different combinations of transcription factors: IL-2 expression is highly dependent on NF-AT and Oct/OAP binding^{11,26}, whereas IL-8 depends on NF- κ B and other factors but not on NF-AT^{25,27}. Cells were transfected with luciferase reporter constructs driven by the IL-2 or IL-8 promoters and were stimulated for 3 h with 50 nM PdBu and constant $[\text{Ca}^{2+}]_i$ elevation. Both genes appear to be highly Ca^{2+} -sensitive, with IL-8 being more sensitive than IL-2 (Fig. 3c). The two genes show different sensitivities to $[\text{Ca}^{2+}]_i$ oscillation frequency (Fig. 3d): IL-2 is not expressed at periods ≥ 400 s, whereas IL-8 retains $\sim 20\%$ of its maximal activity under these conditions. The behaviour of IL-2 resembles that of NF-AT and Oct/OAP, whereas the behaviour of IL-8 is similar to that of NF- κ B. These results were supported by measurements of endogenous IL-8 expression in Jurkat cells treated with monensin to prevent secretion²⁸ (IL-2 expression after 3 h stimulation was too low to measure accurately). FACS analysis of anti-IL-8-stained cells indicated that the frequency dependence of IL-8 expression was similar to that of the reporter gene. Together, these results indicate

that oscillations of different frequencies can lead to the expression of different sets of genes, presumably as a consequence of their effects on the underlying transcription factors.

Our results demonstrate two important properties of nuclear signalling by $[\text{Ca}^{2+}]_i$ oscillations. First, oscillations enhance signalling efficiency specifically at low levels of stimulation. This effect arises from the highly nonlinear dependence of transcription on $[\text{Ca}^{2+}]_i$, so that oscillations periodically exceed the threshold for activation whereas a small constant $[\text{Ca}^{2+}]_i$ rise of the same average magnitude does not. The tendency of $[\text{Ca}^{2+}]_i$ to oscillate at low receptor occupancy in many cells may thus optimize sensitivity to weak external stimuli. Second, oscillations confer specificity on an otherwise highly pleiotropic Ca^{2+} signal. By differentially controlling the activation of distinct sets of transcription factors and the expression of different genes, oscillation frequency may direct cells along specific developmental pathways^{29,30}. Frequency-dependent gene expression is likely to be widespread, as Ca^{2+} -dependent transcription factors like NF- κ B and NF-AT are present in a great variety of cells and oscillations can occur with periods of seconds to hours^{1,29}. □

Methods

Cells and solutions. All experiments were done on the NF-ATZ-DIPA clone of Jurkat T cells, stably transfected with an NF-AT-*lacZ* reporter gene and selected for high *lacZ* inducibility¹³. Experiments were done in RPMI 1640 without




Figure 2 $[\text{Ca}^{2+}]_i$ oscillations enhance the calcium sensitivity of NF-AT-dependent transcription at low levels of stimulation. **a**, NF-AT-dependent *lacZ* expression in cells stimulated with a low level of $[\text{Ca}^{2+}]_i$. Left, superimposed $[\text{Ca}^{2+}]_i$ recordings from two cell populations (200–300 cells each) stimulated for 3 h with oscillating $[\text{Ca}^{2+}]_i$ (227 nM average) or constant $[\text{Ca}^{2+}]_i$ (225 nM average). Right, flow cytometry analysis of β -gal activity in the same experiments show that oscillating $[\text{Ca}^{2+}]_i$ activates 39% of the cells while constant $[\text{Ca}^{2+}]_i$ activates only 19%. The vertical gate excludes 97% of the unstimulated cells. **b**, NF-AT-dependent expression at a high level of $[\text{Ca}^{2+}]_i$. Left, stimulation with oscillating $[\text{Ca}^{2+}]_i$ (340 nM average) or constant $[\text{Ca}^{2+}]_i$ (359 nM average). Right, oscillating and constant $[\text{Ca}^{2+}]_i$ activate 33 and 39% of the cells, respectively. **c**, Percentage of *lacZ*⁺ cells induced by various concentrations of oscillating (○) or constant (●) $[\text{Ca}^{2+}]_i$. **d**, Ratio of the number of β -gal⁺ cells activated by oscillating as compared with constant $[\text{Ca}^{2+}]_i$, as a function of average $[\text{Ca}^{2+}]_i$, calculated from the data in **c**. All experiments were done in the presence of 50 nM PdBu.




Figure 3 Effects of $[\text{Ca}^{2+}]_i$ and oscillation frequency on NF-AT-, Oct/OAP-, NF- κ B-, IL-2- and IL-8-luciferase reporter genes. **a**, Steady-state Ca^{2+} sensitivities of NF-AT, Oct/OAP and NF- κ B. A plot of the Hill equation is superimposed, with a Hill coefficient (n_H) of 4.7 and a K_d of 270 nM. **b**, Oscillation frequency dependences of NF-AT (□), Oct/OAP (Δ) or NF- κ B (○) reporter genes. **c**, Steady-state Ca^{2+} sensitivities of IL-2 (○) and IL-8 (Δ) reporter genes. Fits to the Hill equation are superimposed (for IL-2, $n_H = 2.7$ and $K_d = 346$ nM; for IL-8, $n_H = 3.6$ and $K_d = 166$ nM). **d**, Frequency dependences of IL-2 (○) and IL-8 (Δ) reporter genes. Error bars represent s.e.m. of triplicate measurements in 3–5 experiments; where they are not visible they are smaller than the symbols. All experiments were done in the presence of 50 nM PdBu.

phenol red or riboflavins and supplemented with 25 mM HEPES and 0.2% BSA. Solutions were maintained at 4 °C until immediately before each experiment, when they were warmed to 37 °C and supplemented with 50 nM PdBU and either 0.5 mM EGTA (~50 μM free Ca²⁺) or 1 mM Ca²⁺ (1.5 mM total Ca²⁺).

Transfection of the reporter genes. NF-AT, NF-κB, Oct-1/OAP and IL-2 reporter constructs were provided by J. Goldberg and G. Crabtree and were derived from constructs described previously¹⁶. Multimeric copies of NFAT (3 copies), NF-κB (4) or Oct/Oap (4) binding sites, linked to a minimal (non-inducible) IL-2 promoter (-74 to +47), were inserted between the *Sma*I and *Hind*III sites in the multiple cloning region of the pGL-3 luciferase reporter vector (Promega). The construct with the intact IL-2 promoter region was made by inserting 371 bp of the IL-2 initiation region (-324 to +47) into the pGL-3 vector. The IL-8 reporter construct was provided by K. P. LeClair²⁷.

Transfection of 10⁷ cells was by electroporation with 10 μg of the reporter vector, 1 μg of a vector containing large-T antigen and 2 μg of a vector encoding the transmembrane and extracellular domains of CD8. Large-T antigen was used to increase the number of copies of the reporter construct in each cell, and the CD8 construct was used to determine the transfection efficiency. Transfection efficiencies were 30–40%; viability following centrifugation to remove cells killed during electroporation was 85–95%. Experiments were conducted 24–48 h after transfection when the expression of reporter genes was maximal.

Ca²⁺ clamp and Ca²⁺ measurements. Transfected cells were loaded with 2 μM Fura-PE3/AM (Teflabs) for 1 h at 37 °C in loading medium (RPMI 1640, 25 mM HEPES, 2% fetal bovine serum), washed, and incubated for another hour to allow complete de-esterification of the dye. Loaded cells (2–3 × 10⁵) were allowed to adhere to a polylysine-coated laminar flow chamber and were placed on the heated stage of a Zeiss Axiovert 35 inverted microscope. The laminar flow chamber (60 μl volume) was connected to two heated reservoirs containing 1.5 mM and 0 mM Ca²⁺ solutions, and pressurized with a mixture of 95% air and 5% CO₂. At the start of each experiment, cells were treated with 1 μM thapsigargin in 0 mM Ca²⁺ solution for 5 min to deplete internal Ca²⁺ stores and irreversibly activate CRAC channels. A computer-controlled solenoid valve (General Valve) was used to switch rapidly between the Ca²⁺-containing and Ca²⁺-free solutions flowing into the chamber and over the cells. The solution in the chamber was fully exchanged about once per second and was maintained at 37 °C. Cells were stimulated for 3 h while [Ca²⁺]_i was measured every 5–10 s by video microscopy as described¹⁰. Fura-PE3 was calibrated on the microscope in a microcuvette using solutions containing 1 mM EGTA and 10 mM Ca²⁺ according to ref. 10.

Reporter gene assays. Following stimulation cells were washed from the chamber, lysed by freeze/thawing and subjected to a luciferase assay using standard methods. All measurements were done in triplicate and normalized to the total number of cells determined using a Coulter Counter (Coulter Electronics). β-Galactosidase (β-gal) expression was measured by flow cytometry using the FACS-Gal protocol¹³. Steady-state [Ca²⁺]_i dependence of luciferase reporter genes (Fig. 3) was determined after 3 h stimulation with 1 μM thapsigargin, 50 nM PdBU and variable extracellular [Ca²⁺]. Luciferase activity was normalized to values between 0 and 100%, given by the responses to TG + PdBU in medium containing 0.5 mM EGTA or 1.5 mM Ca²⁺, respectively.

Received 16 December 1997; accepted 12 February 1998.

- Berridge, M. J. & Galione, A. Cytosolic calcium oscillators. *FASEB J.* **2**, 3074–3082 (1988).
- Tsien, R. W. & Tsien, R. Y. Calcium channels, stores, and oscillations. *Annu. Rev. Cell Biol.* **6**, 715–760 (1990).
- Fewtrell, C. Ca²⁺ oscillations in non-excitable cells. *Annu. Rev. Physiol.* **55**, 427–454 (1993).
- Thomas, A. P., Bird, G. S. J., Hajnóczky, G., Robb-Gaspers, L. D. & Putney, J. W. Jr Spatial and temporal aspects of cellular calcium signaling. *FASEB J.* **10**, 1505–1517 (1996).
- Negulescu, P. A., Shastri, N. & Cahalan, M. D. Intracellular calcium dependence of gene expression in single T lymphocytes. *Proc. Natl Acad. Sci. USA* **91**, 2873–2877 (1994).
- Dolmetsch, R. E. & Lewis, R. S. in *Imaging Living Cells: A Laboratory Manual* (eds Konnerth, A., Lanni, F. & Yuste, R.) (CSHL Press, Cold Spring Harbor, in the press).
- Lewis, R. S. & Cahalan, M. D. Potassium and calcium channels in lymphocytes. *Annu. Rev. Immunol.* **13**, 623–653 (1995).
- Zweifach, A. & Lewis, R. S. Mitogen-regulated Ca²⁺ current of T lymphocytes is activated by depletion of intracellular Ca²⁺ stores. *Proc. Natl Acad. Sci. USA* **90**, 6295–6299 (1993).
- Lewis, R. S. & Cahalan, M. D. Mitogen-induced oscillations of cytosolic Ca²⁺ and transmembrane Ca²⁺ current in human leukemic T cells. *Cell Reg.* **1**, 99–112 (1989).
- Dolmetsch, R. E. & Lewis, R. S. Signaling between intracellular Ca²⁺ stores and depletion-activated Ca²⁺ channels generates [Ca²⁺]_i oscillations in T lymphocytes. *J. Gen. Physiol.* **103**, 365–388 (1994).

- Crabtree, G. R. & Clipstone, N. A. Signal transmission between the plasma membrane and nucleus of T lymphocytes. *Annu. Rev. Biochem.* **63**, 1045–1083 (1994).
- Rao, A. NF-ATp: a transcription factor required for the coordinate induction of several cytokine genes. *Immunol. Today* **15**, 274–281 (1994).
- Fiering, S. et al. Single cell assay of a transcription factor reveals a threshold in transcription activated by signals emanating from the T-cell antigen receptor. *Genes Dev.* **4**, 1823–1834 (1990).
- Donnadieu, E. et al. Imaging early steps of human T cell activation by antigen-presenting cells. *J. Immunol.* **148**, 2643–2653 (1992).
- Frantz, B. et al. Calcineurin acts in synergy with PMA to inactivate IκB/MAD3, an inhibitor of NF-κB. *EMBO J.* **13**, 861–870 (1994).
- Mattila, P. S. et al. The actions of cyclosporin A and FK506 suggest a novel step in the activation of T lymphocytes. *EMBO J.* **9**, 4425–4433 (1990).
- Baeuerle, P. A. & Henkel, T. Function and activation of NF-κB in the immune system. *Annu. Rev. Immunol.* **12**, 141–179 (1994).
- Clipstone, N. A. & Crabtree, G. R. Identification of calcineurin as a key signalling enzyme in T-lymphocyte activation. *Nature* **357**, 695–697 (1992).
- Liu, J. et al. Calcineurin is a common target of cyclophilin-cyclosporin A and FKBP-FK506 complexes. *Cell* **66**, 807–815 (1991).
- Fanger, C. M., Hoth, M., Crabtree, G. R. & Lewis, R. S. Characterization of T cell mutants with defects in capacitative calcium entry: Genetic evidence for the physiological roles of CRAC channels. *J. Cell Biol.* **131**, 655–667 (1995).
- Dolmetsch, R. E., Lewis, R. S., Goodnow, C. C. & Healy, J. I. Differential activation of transcription factors induced by Ca²⁺ response amplitude and duration. *Nature* **386**, 855–858 (1993).
- Li, W.-h., Llopis, J., Whitney, M., Zlokarnik, G. & Tsien, R. Y. Cell-permeant caged InsP₃ ester shows that Ca²⁺ spike frequency can optimize gene expression. *Nature* **392**, 936–941 (1998).
- Shibasaki, F., Price, E., Milan, D. & McKeon, F. Role of kinases and the phosphatase calcineurin in the nuclear shuttling of transcription factor NF-AT4. *Nature* **382**, 370–373 (1996).
- Durand, D. B., Bush, M. R., Morgan, J. G., Weiss, A. & Crabtree, G. R. A 275 base pair fragment at the 5' end of the interleukin 2 gene enhances expression from a heterologous promoter in response to signals from the T cell antigen receptor. *J. Exp. Med.* **165**, 395–407 (1987).
- Okamoto, S. et al. The interleukin-8 AP-1 and κB-like sites are genetic end targets of FK506-sensitive pathway accompanied by calcium mobilization. *J. Biol. Chem.* **269**, 8582–8589 (1994).
- Durand, D. B. et al. Characterization of antigen receptor response elements within the interleukin-2 enhancer. *Mol. Cell. Biol.* **8**, 1715–1724 (1988).
- Wechsler, A. S., Gordon, M. C., Dendorfer, U. & LeClair, K. P. Induction of IL-8 expression in T cells uses the CD28 costimulatory pathway. *J. Immunol.* **153**, 2515–2523 (1994).
- Prussin, C. & Metcalfe, D. D. Detection of intracytoplasmic cytokine using flow cytometry and directly conjugated anti-cytokine antibodies. *J. Immunol. Meth.* **188**, 117–128 (1995).
- Gu, X. & Spitzer, N. C. Distinct aspects of neuronal differentiation encoded by frequency of spontaneous Ca²⁺ transients. *Nature* **375**, 784–787 (1995).
- Fields, R. D., Eshete, F., Stevens, B. & Itoh, K. Action potential-dependent regulation of gene expression: temporal specificity in Ca²⁺, cAMP-responsive element binding proteins, and mitogen-activated protein kinase signaling. *J. Neurosci.* **17**, 7252–7266 (1997).

Acknowledgements. We thank J. Goldberg for NF-AT-, Oct/OAP- and NF-κB-luciferase constructs; K. LeClair and G. Crabtree for IL-8 and IL-2 luciferase constructs; C. Fanger for creating the large T antigen and CD8 constructs; G. Crabtree and L. Herzenberg for use of the luminometer and electroporator; S. Finkbeiner, Y. Gotoh and M. Greenberg for comments on the manuscript; and J. Walton for help in preparing constructs. This work was supported by a grant from the NIH (R.S.L.) and by a predoctoral fellowship from the American Heart Association, California Affiliate (R.E.D.).

Correspondence and requests for materials should be addressed to R.S.L. (e-mail: rslweis@leland.stanford.edu).

Cell-permeant caged InsP₃ ester shows that Ca²⁺ spike frequency can optimize gene expression

Wen-hong Li^{†‡}, Juan Llopis^{*}, Michael Whitney[§], Gregor Zlokarnik[§] & Roger Y. Tsien^{††}

Departments of ^{*} Pharmacology and [†] Chemistry & Biochemistry, and [‡] Howard Hughes Medical Institute, University of California, San Diego, 9500 Gilman Drive, La Jolla, California 92093-0647, USA
[§] Aurora Biosciences Corporation, 11010 Torreyana Road, San Diego, California 92121, USA

Inositol 1,4,5-trisphosphate (InsP₃) releases calcium from intracellular stores and triggers complex waves and oscillations in levels of cytosolic free calcium^{1–5}. To determine which longer-term responses are controlled by oscillations in InsP₃ and cytosolic free calcium, it would be useful to deliver exogenous InsP₃, under spatial and temporal control, into populations of unpermeabilized cells. Here we report the 15-step synthesis of a

[‡] Present address: Division of Biology 139-74, Beckman Institute, California Institute of Technology, Pasadena, California 91125, USA.