


Properties of Sound



- What is sound?
- Wavelength, period, frequency
- Interference and the linearity of sound
- Sound attenuation
 - Spreading loss
 - Acoustic impedance
 - Scattering
- Reading in Chap 2
 - box 2.3 but not 2.1 or 2.2




How does a cicada sing?



(B)




Sound is a pressure wave



- Period (τ) = the time between successive waveform peaks
- Frequency (in cycles/sec or Hertz) = $1/\text{period}$,
- Thus $f = 1/\tau$

Wavelength depends on speed of sound



- Wavelength of a sound is the distance traveled in one cycle (or period)
- Wavelength = period • speed of sound in medium (c)
- Thus wavelength = $c / \text{frequency}$

Acoustical property comparisons

<u>Property</u>	<u>Air</u>	<u>Water</u>	<u>Rock</u>
Speed (m/s)	340	1500	2000-5000
Density (g/cm ³)	0.001	1	2-3

Wavelength problem

Which sound has a shorter wavelength:
1 kHz in air or 3 kHz in water?


Wavelength = speed of sound / frequency

Air: $340 \text{ m/s} / 1000 \text{ cycle/s} = 0.34 \text{ m/cycle}$

Water: $1500 \text{ m/s} / 3000 \text{ cycle/s} = 0.5 \text{ m/cycle}$

Therefore, the answer is 1 kHz in air

Frequencies and wavelengths in air




Pressure vs particle movements



Close to a sound source, pressure and particle displacement are out of phase, but one wavelength from source they are in phase

Near field vs far field

- Near field
 - molecular displacements $>$ pressure differences
 - About 1-2 wavelengths from source
- Far field
 - pressure displacement $>$ molecular displacements
 - travels as a wave away from source




The Doppler shift

- When the sound source is moving, the frequency of the sound will be altered. This is known as the Doppler shift
- Approaching sounds are higher in frequency
- Receding sounds are lower in frequency
- See demonstration at
 - <http://www.walter-fendt.de/ph14e/dopplereff.htm>



Sound linearity and interference

- Sound pressure waves combine additively
- Waves that start together are in phase
 - Sounds in phase increase in amplitude (positive interference)
 - Sounds out of phase cancel each other out (negative interference)
 - Sounds partially out of phase create varying amplitudes (beats)

Positive interference




+




Summed Waveform


Negative Interference




+




Interference produces Beats



+




Summed Waveform



Amplitude measures

Peak Peak-to-peak Root-mean-squared (RMS)



Amplitude scale

- Sound pressure is measured in decibels (dB) on a \log_{10} scale relative to a reference level
- $\text{dB} = 20 \log_{10} P_1/P_r$ where P_r is a reference pressure level
- A common reference pressure level is the threshold of human hearing at 1 kHz, referred to as sound pressure level (SPL)
- A sound with twice the SPL is 6 dB louder
 - i.e. $20 \log_{10} (2) = 20(0.3) = 6$

Sample sound pressure levels

- soft whisper 20 dB
- nearby songbird, office hum 50 dB
- barking dog 70 dB
- roaring lion , heavy truck 90 dB
- echolocating big brown bat 100 dB
- jet take-off 120 dB

Amplitude problems

- If sound A has 10 times the SPL of sound B, how much louder in dB is A than B?

$$\text{dB} = 20 \log_{10} 10 = 20 \text{ dB louder}$$

- If sound A is 100 db and sound B is 80 db, how much louder is A than B?

20 db


- If an 80 db sound is combined with a 40 db sound, how loud is the sound (approximately)?

80 db

Sound attenuation

- Spherical spreading
- Absorption
 - Temperature and humidity effects
- Reflective scattering
 - Due to impedance differences


Spherical spreading




- Loss in sound intensity follows the inverse square law
 - pressure drops in half for each doubling of distance, i.e. for each doubling of distance sound is 6 dB less

Atmospheric attenuation

Increases with
square of frequency



Nonlinear with humidity




Acoustic impedance

- The degree to which a medium is compressible by sound pressure
- Acoustic impedance = speed of sound • density of medium
- Transmission between media with different impedances is difficult
 - Sounds reflect off animals in air, but can pass through them in water
- Dictates efficiency of sound production and reception by organisms

Acoustical property comparisons

<u>Property</u>	<u>Air</u>	<u>Water</u>	<u>Rock</u>
Speed (m/s)	340	1500	2000-5000
Density (g/cm ³)	0.001	1	2-3
Impedance(rayls)	30	1.5 x 10⁵	4-5 x 10⁵

Reflection and refraction




- If medium 2 is more dense than medium 1
 - Most of the energy in a sound wave will be reflected from the boundary between media
 - If the incidence angle exceeds a threshold (see book), some energy will be refracted into media 2, with a reduced angle
- Reflection depends on size of media 2 object


Reflective Scattering

- Type of scattering depends on ratio of wavelength and reflecting object
 - Rayleigh scattering (object \ll wavelength)
 - sound scattered equally in all directions
 - Diffractive or Mie scattering (object = wavelength)
 - both a reflected and diffracted wave
 - Simple scattering (object $>$ wavelength)
 - single reflected wave

Scattering and wavelengths



Diffractive scattering



- Part of sound wave is diffracted around object (creeping wave)
- Reflected wave is out of phase with creeping wave.
- Can cause negative interference